

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS
-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------

All races 3/, both sexes	*	*	*	*	*	*	*	*
Male	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*
White, both sexes	*	*	*	*	*	*	*	*
Male	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*
Black, both sexes	*	*	*	*	*	*	*	*
Male	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*

All races 3/, both sexes	0.5	*	*	*	*	*	*	0.1	0.3
Male	0.4	*	*	*	*	*	*	*	0.3
Female	0.6	*	*	*	*	*	*	*	0.2
White, both sexes	0.5	*	*	*	*	*	*	0.1	0.3
Male	0.4	*	*	*	*	*	*	*	0.3
Female	0.6	*	*	*	*	*	*	*	0.3
Black, both sexes	0.2	*	*	*	*	*	*	*	*
Male	0.2	*	*	*	*	*	*	*	*
Female	0.3	*	*	*	*	*	*	*	*

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS
----------------	--------------------	--------------	---------------	----------------	----------------	----------------	----------------	----------------

Measles (B05)

All races 3/, both sexes	*	*	*	*	*	*	*	*
Male	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*
White, both sexes	*	*	*	*	*	*	*	*
Male	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*
Black, both sexes	*	*	*	*	*	*	*	*
Male	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*

Viral hepatitis (B15-B19)

All races 3/, both sexes	1.9	*	*	*	*	0.2	1.7	5.1	3.8
Male	2.5	*	*	*	*	0.2	2.4	7.8	5.0
Female	1.4	*	*	*	*	0.2	1.1	2.5	2.6
White, both sexes	1.9	*	*	*	*	0.2	1.7	4.7	3.2
Male	2.4	*	*	*	*	0.2	2.4	7.3	4.3
Female	1.3	*	*	*	*	0.1	1.0	2.1	2.2
Black, both sexes	2.2	*	*	*	*	0.4	2.1	8.1	5.9
Male	3.0	*	*	*	*	*	2.9	12.8	8.4
Female	1.6	*	*	*	*	*	1.3	4.2	4.0

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS
----------------	--------------------	--------------	---------------	----------------	----------------	----------------	----------------	----------------

Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09,B25-B49,B55-B99)

All races 3/, both sexes	2.0	4.3	0.5	0.2	0.3	0.4	0.8	1.6	2.9
Male	2.1	4.8	0.5	0.2	0.3	0.4	1.0	2.0	3.3
Female	1.9	3.9	0.4	0.2	0.2	0.4	0.6	1.2	2.6
White, both sexes	2.0	3.3	0.4	0.2	0.2	0.3	0.7	1.4	2.7
Male	2.1	3.8	0.4	0.2	0.3	0.3	0.8	1.8	3.1
Female	1.9	2.8	0.4	0.2	0.2	0.2	0.5	1.1	2.3
Black, both sexes	2.3	10.5	*	*	0.4	1.0	1.7	3.1	4.9
Male	2.6	10.5	*	*	*	0.9	2.2	4.0	5.9
Female	2.1	10.5	*	*	*	1.0	1.2	2.3	4.2

Malignant neoplasms (C00-C97)

All races 3/, both sexes	200.9	2.4	2.8	2.6	4.5	10.5	36.8	129.3	371.3
Male	212.5	2.5	3.0	2.8	5.2	10.0	32.9	133.3	423.5
Female	189.9	2.2	2.5	2.3	3.7	11.0	40.6	125.5	323.6
White, both sexes	212.2	2.5	2.7	2.6	4.4	10.3	34.6	122.9	362.8
Male	222.5	2.7	3.1	2.9	5.1	10.0	31.0	125.0	406.8
Female	202.2	2.2	2.4	2.4	3.6	10.6	38.2	120.9	321.7
Black, both sexes	175.5	*	3.1	2.4	5.2	12.4	54.5	197.3	502.4
Male	195.6	*	3.0	2.7	6.1	10.5	48.8	223.2	654.3
Female	157.2	*	3.3	2.0	4.2	14.1	59.5	175.7	386.6

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS
----------------	--------------------	--------------	---------------	----------------	----------------	----------------	----------------	----------------

Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)

All races 3/, both sexes	2.7	*	*	*	0.1	0.1	0.7	2.9	6.8
Male	3.7	*	*	*	*	0.2	1.0	4.5	11.1
Female	1.8	*	*	*	*	0.1	0.3	1.3	3.0
White, both sexes	2.7	*	*	*	*	0.1	0.5	2.4	6.1
Male	3.5	*	*	*	*	0.1	0.7	3.7	9.7
Female	1.9	*	*	*	*	*	0.3	1.2	2.7
Black, both sexes	3.1	*	*	*	*	*	1.1	6.1	13.7
Male	5.0	*	*	*	*	*	1.8	11.0	25.4
Female	1.5	*	*	*	*	*	*	2.0	4.8

Malignant neoplasm of esophagus (C15)

All races 3/, both sexes	4.4	*	*	*	*	0.1	0.7	3.6	10.9
Male	6.9	*	*	*	*	0.2	1.2	6.4	19.0
Female	2.1	*	*	*	*	*	0.1	1.0	3.5
White, both sexes	4.6	*	*	*	*	0.1	0.7	3.4	10.1
Male	7.2	*	*	*	*	0.2	1.2	6.0	17.8
Female	2.1	*	*	*	*	*	0.1	0.8	3.0
Black, both sexes	4.7	*	*	*	*	*	0.8	6.8	20.0
Male	7.0	*	*	*	*	*	1.2	11.7	35.5
Female	2.6	*	*	*	*	*	*	2.7	8.2

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS
----------------	--------------------	--------------	---------------	----------------	----------------	----------------	----------------	----------------

Malignant neoplasm of stomach (C16)

All races 3/, both sexes	4.6	*	*	*	*	0.3	1.1	3.0	7.5
Male	5.4	*	*	*	*	0.4	1.3	3.9	10.7
Female	3.8	*	*	*	*	0.3	0.9	2.1	4.6
White, both sexes	4.3	*	*	*	*	0.3	0.9	2.5	6.6
Male	5.1	*	*	*	*	0.3	1.0	3.2	9.4
Female	3.5	*	*	*	*	0.2	0.7	1.7	3.9
Black, both sexes	6.2	*	*	*	*	0.7	2.1	5.8	14.0
Male	7.3	*	*	*	*	0.8	2.9	8.3	21.6
Female	5.2	*	*	*	*	*	1.3	3.8	8.3

Malignant neoplasms of colon, rectum and anus (C18-C21)

All races 3/, both sexes	20.9	*	*	*	0.1	0.7	3.0	11.4	33.1
Male	21.1	*	*	*	0.1	0.8	3.4	13.0	40.3
Female	20.6	*	*	*	*	0.6	2.7	9.9	26.6
White, both sexes	21.9	*	*	*	0.1	0.7	2.8	10.5	31.5
Male	22.0	*	*	*	*	0.8	3.1	12.1	38.1
Female	21.7	*	*	*	*	0.6	2.4	9.0	25.3
Black, both sexes	19.5	*	*	*	*	1.0	5.1	19.9	50.8
Male	19.8	*	*	*	*	1.0	6.0	22.9	65.3
Female	19.1	*	*	*	*	1.0	4.4	17.4	39.8

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS
----------------	--------------------	--------------	---------------	----------------	----------------	----------------	----------------	----------------

Malignant neoplasms of liver and intrahepatic bile ducts (C22)

All races 3/, both sexes	4.7	*	0.2	0.1	0.1	0.2	1.0	4.5	9.4
Male	6.1	*	*	*	0.1	0.3	1.3	7.3	14.1
Female	3.4	*	*	*	*	0.1	0.6	1.9	5.2
White, both sexes	4.6	*	*	*	0.1	0.1	0.7	3.7	8.2
Male	5.8	*	*	*	*	0.2	1.0	5.8	11.9
Female	3.4	*	*	*	*	*	0.4	1.7	4.7
Black, both sexes	4.5	*	*	*	*	0.4	1.8	8.8	14.3
Male	6.3	*	*	*	*	*	2.5	15.4	24.5
Female	2.9	*	*	*	*	*	1.2	3.3	6.4

Malignant neoplasm of pancreas (C25)

All races 3/, both sexes	10.7	*	*	*	*	0.1	1.2	6.4	19.6
Male	10.6	*	*	*	*	0.2	1.5	8.2	24.0
Female	10.7	*	*	*	*	0.1	1.0	4.6	15.6
White, both sexes	11.2	*	*	*	*	0.2	1.2	6.2	19.0
Male	11.1	*	*	*	*	0.2	1.4	7.9	23.3
Female	11.2	*	*	*	*	*	0.9	4.4	15.1
Black, both sexes	9.7	*	*	*	*	*	1.7	9.3	27.9
Male	9.3	*	*	*	*	*	2.0	12.3	35.7
Female	10.0	*	*	*	*	*	1.4	6.7	21.9

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS
----------------	--------------------	--------------	---------------	----------------	----------------	----------------	----------------	----------------

Malignant neoplasm of larynx (C32)

All races 3/, both sexes	1.4	*	*	*	*	*	0.1	1.1	3.6
Male	2.3	*	*	*	*	*	0.2	1.9	6.3
Female	0.6	*	*	*	*	*	*	0.5	1.2
White, both sexes	1.4	*	*	*	*	*	0.1	1.0	3.1
Male	2.2	*	*	*	*	*	0.2	1.5	5.3
Female	0.6	*	*	*	*	*	*	0.4	1.1
Black, both sexes	2.0	*	*	*	*	*	*	3.0	9.1
Male	3.4	*	*	*	*	*	*	5.3	17.9
Female	0.8	*	*	*	*	*	*	1.0	2.3

Malignant neoplasms of trachea, bronchus and lung (C33-C34)

All races 3/, both sexes	56.5	*	*	*	0.1	0.5	6.1	32.1	124.0
Male	67.2	*	*	*	*	0.5	6.9	39.2	156.9
Female	46.3	*	*	*	*	0.5	5.4	25.2	94.0
White, both sexes	60.5	*	*	*	*	0.5	5.7	30.5	123.7
Male	70.8	*	*	*	*	0.5	6.1	36.1	152.6
Female	50.6	*	*	*	*	0.5	5.2	24.9	96.7
Black, both sexes	45.5	*	*	*	*	0.6	10.2	51.6	154.5
Male	60.5	*	*	*	*	*	13.5	73.7	233.5
Female	31.9	*	*	*	*	*	7.3	33.2	94.2

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS
----------------	--------------------	--------------	---------------	----------------	----------------	----------------	----------------	----------------

Malignant melanoma of skin (C43)

All races 3/, both sexes	2.7	*	*	*	0.1	0.5	1.4	3.2	5.4
Male	3.4	*	*	*	0.1	0.6	1.6	4.2	7.6
Female	2.0	*	*	*	0.1	0.4	1.2	2.3	3.5
White, both sexes	3.2	*	*	*	0.1	0.6	1.6	3.8	6.2
Male	4.1	*	*	*	0.1	0.7	1.9	4.9	8.6
Female	2.4	*	*	*	*	0.5	1.4	2.6	4.0
Black, both sexes	0.4	*	*	*	*	*	*	*	0.8
Male	0.4	*	*	*	*	*	*	*	*
Female	0.4	*	*	*	*	*	*	*	*

Malignant neoplasm of breast (C50)

All races 3/, both sexes	15.4	*	*	*	*	1.2	6.3	17.2	31.6
Male	0.3	*	*	*	*	*	*	0.3	0.7
Female	29.8	*	*	*	*	2.5	12.5	33.4	59.7
White, both sexes	16.0	*	*	*	*	1.1	5.7	16.0	30.5
Male	0.3	*	*	*	*	*	*	0.2	0.7
Female	31.1	*	*	*	*	2.2	11.4	31.4	58.3
Black, both sexes	15.4	*	*	*	*	2.3	11.3	28.5	46.0
Male	0.4	*	*	*	*	*	*	*	*
Female	28.9	*	*	*	*	4.4	21.1	52.0	80.0

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS
----------------	--------------------	--------------	---------------	----------------	----------------	----------------	----------------	----------------

Malignant neoplasm of cervix uteri (C53)

All races 3/, both sexes	1.5	*	*	*	*	0.6	1.5	2.5	3.1
Female	3.0	*	*	*	*	1.2	3.1	4.8	6.0
White, both sexes	1.4	*	*	*	*	0.6	1.4	2.2	2.7
Female	2.7	*	*	*	*	1.2	2.8	4.3	5.2
Black, both sexes	2.5	*	*	*	*	1.0	2.6	4.4	6.3
Female	4.7	*	*	*	*	1.9	5.0	8.0	11.2

Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)

All races 3/, both sexes	2.4	*	*	*	*	0.1	0.3	1.4	4.7
Female	4.7	*	*	*	*	0.1	0.6	2.7	8.9
White, both sexes	2.4	*	*	*	*	0.1	0.3	1.3	4.3
Female	4.7	*	*	*	*	0.1	0.6	2.7	8.3
Black, both sexes	3.0	*	*	*	*	*	0.4	1.6	8.4
Female	5.7	*	*	*	*	*	0.7	3.0	14.8

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS
----------------	--------------------	--------------	---------------	----------------	----------------	----------------	----------------	----------------

Malignant neoplasm of ovary (C56)

All races 3/, both sexes	5.1	*	*	*	0.1	0.3	1.1	4.3	10.2
Female	10.0	*	*	*	0.1	0.5	2.1	8.3	19.5
White, both sexes	5.6	*	*	*	*	0.3	1.1	4.4	10.6
Female	11.1	*	*	*	*	0.6	2.2	8.7	20.5
Black, both sexes	3.0	*	*	*	*	*	1.1	3.4	8.5
Female	5.8	*	*	*	*	*	2.0	6.2	15.0

Malignant neoplasm of prostate (C61)

All races 3/, both sexes	11.3	*	*	*	*	*	0.1	1.0	8.0
Male	23.1	*	*	*	*	*	0.1	2.0	16.9
White, both sexes	11.2	*	*	*	*	*	*	0.8	6.8
Male	22.8	*	*	*	*	*	*	1.7	14.1
Black, both sexes	15.1	*	*	*	*	*	*	2.5	20.6
Male	31.9	*	*	*	*	*	*	5.4	47.6

UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

OF DISEASES, 1992), RACE, AND SEX	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS
Malignant neoplasms of kidney and renal pelvis (C64-C65)									
All races 3/, both sexes	4.3	*	*	0.1	0.1	0.1	0.7	3.3	9.2
Male	5.4	*	*	*	*	0.1	1.0	4.8	13.4
Female	3.2	*	*	*	*	*	0.5	1.9	5.3
White, both sexes	4.6	*	*	*	*	0.1	0.7	3.4	9.4
Male	5.8	*	*	*	*	*	1.0	4.9	13.6
Female	3.5	*	*	*	*	*	0.4	1.9	5.4
Black, both sexes	3.0	*	*	*	*	*	0.9	3.5	8.9
Male	3.8	*	*	*	*	*	1.0	5.1	13.7
Female	2.2	*	*	*	*	*	0.8	2.2	5.2

Malignant neoplasm of bladder (C67)

[illegible]

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS
----------------	--------------------	--------------	---------------	----------------	----------------	----------------	----------------	----------------

Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)

All races 3/, both sexes	4.6	0.5	0.7	0.8	0.6	1.1	2.4	5.2	9.7
Male	5.2	*	0.7	0.9	0.7	1.4	3.2	6.3	12.1
Female	4.0	*	0.6	0.7	0.5	0.8	1.7	4.1	7.6
White, both sexes	5.2	*	0.7	0.8	0.6	1.2	2.6	5.7	10.5
Male	5.8	*	0.7	0.9	0.7	1.5	3.4	6.8	12.9
Female	4.5	*	0.6	0.7	0.5	0.9	1.8	4.5	8.2
Black, both sexes	2.2	*	*	0.8	0.6	0.7	1.6	3.0	5.8
Male	2.5	*	*	1.0	0.7	1.1	2.0	3.9	7.4
Female	1.9	*	*	0.7	*	*	1.2	2.2	4.5

Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)

All races 3/, both sexes	20.5	1.0	1.0	0.9	1.9	2.5	4.1	10.8	30.4
Male	22.2	1.1	1.0	1.1	2.3	2.9	4.8	13.2	36.8
Female	18.9	*	0.9	0.8	1.4	2.0	3.4	8.6	24.6
White, both sexes	22.3	1.2	1.0	1.0	1.8	2.4	4.0	10.5	30.4
Male	24.1	*	1.0	1.1	2.2	2.8	4.8	12.8	36.6
Female	20.5	*	1.0	0.8	1.3	1.9	3.2	8.3	24.5
Black, both sexes	14.4	*	0.9	0.7	2.3	3.1	5.4	15.1	36.3
Male	15.2	*	*	0.8	2.7	3.7	5.7	18.7	46.5
Female	13.6	*	*	*	1.8	2.5	5.1	12.1	28.5

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS
----------------	--------------------	--------------	---------------	----------------	----------------	----------------	----------------	----------------

Hodgkin's disease (C81)

All races 3/, both sexes	0.5	*	*	*	0.2	0.4	0.3	0.4	0.7
Male	0.5	*	*	*	0.2	0.4	0.4	0.5	0.8
Female	0.4	*	*	*	0.2	0.4	0.3	0.3	0.6
White, both sexes	0.5	*	*	*	0.2	0.4	0.3	0.5	0.7
Male	0.6	*	*	*	0.2	0.5	0.4	0.6	0.7
Female	0.5	*	*	*	0.2	0.4	0.3	0.3	0.6
Black, both sexes	0.3	*	*	*	*	*	0.4	*	*
Male	0.3	*	*	*	*	*	*	*	*
Female	0.3	*	*	*	*	*	*	*	*

Non-Hodgkin's lymphoma (C82-C85)

All races 3/, both sexes	8.3	*	*	0.1	0.5	0.8	1.7	4.7	13.2
Male	8.8	*	*	0.1	0.6	0.9	2.2	6.1	16.2
Female	7.8	*	*	*	0.3	0.6	1.2	3.3	10.5
White, both sexes	9.2	*	*	0.1	0.4	0.7	1.7	4.7	13.8
Male	9.7	*	*	0.2	0.6	0.9	2.2	6.1	16.7
Female	8.7	*	*	*	0.2	0.6	1.1	3.4	11.2
Black, both sexes	3.9	*	*	*	0.6	1.1	2.1	5.1	10.7
Male	4.3	*	*	*	*	1.2	2.6	7.2	15.1
Female	3.5	*	*	*	*	0.9	1.7	3.3	7.2

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS
----------------	--------------------	--------------	---------------	----------------	----------------	----------------	----------------	----------------

Leukemia (C91-C95)

All races 3/, both sexes	7.8	1.0	0.9	0.8	1.2	1.3	1.8	3.8	10.1
Male	8.7	1.1	0.9	0.9	1.5	1.5	1.9	4.4	12.7
Female	6.8	*	0.9	0.6	0.9	1.0	1.7	3.3	7.7
White, both sexes	8.5	1.2	1.0	0.8	1.2	1.2	1.8	3.7	10.1
Male	9.5	*	0.9	1.0	1.4	1.4	1.9	4.2	12.7
Female	7.5	*	1.0	0.7	0.9	0.9	1.6	3.2	7.7
Black, both sexes	4.9	*	*	0.6	1.4	1.7	2.1	4.9	10.8
Male	5.5	*	*	0.7	1.9	2.0	2.0	5.7	14.5
Female	4.4	*	*	*	0.9	1.4	2.1	4.3	8.0

Multiple myeloma and immunoproliferative neoplasms (C88,C90)

All races 3/, both sexes	4.0	*	*	*	*	*	0.3	1.9	6.4
Male	4.1	*	*	*	*	*	0.4	2.2	7.1
Female	3.9	*	*	*	*	*	0.3	1.6	5.8
White, both sexes	4.0	*	*	*	*	*	0.3	1.6	5.6
Male	4.2	*	*	*	*	*	0.3	1.8	6.4
Female	3.8	*	*	*	*	*	0.2	1.4	5.0
Black, both sexes	5.2	*	*	*	*	*	0.8	4.6	14.1
Male	5.0	*	*	*	*	*	*	5.2	15.7
Female	5.4	*	*	*	*	*	0.9	4.1	12.8

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS
----------------	--------------------	--------------	---------------	----------------	----------------	----------------	----------------	----------------

Other and unspecified malignant neoplasms of lymphoid, hematopoietic and related tissue (C96)

All races 3/, both sexes	0.0	*	*	*	*	*	*	*
Male	0.0	*	*	*	*	*	*	*
Female	0.0	*	*	*	*	*	*	*
White, both sexes	0.0	*	*	*	*	*	*	*
Male	0.0	*	*	*	*	*	*	*
Female	0.0	*	*	*	*	*	*	*
Black, both sexes	*	*	*	*	*	*	*	*
Male	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*

All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)

All races 3/, both sexes	22.9	0.6	0.9	0.7	1.3	1.9	4.7	14.3	38.8
Male	23.6	*	1.0	0.7	1.6	2.3	5.1	16.3	45.7
Female	22.3	*	0.7	0.6	1.0	1.4	4.2	12.4	32.5
White, both sexes	24.3	*	0.8	0.7	1.3	1.9	4.3	13.4	37.8
Male	24.9	*	1.0	0.7	1.6	2.5	4.7	15.3	43.8
Female	23.8	*	0.6	0.6	1.1	1.4	3.9	11.6	32.3
Black, both sexes	19.0	*	1.1	0.7	1.5	1.7	7.6	22.6	52.5
Male	20.1	*	*	0.8	1.9	1.9	8.6	27.3	71.1
Female	17.9	*	*	*	1.0	1.5	6.7	18.7	38.4

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS
----------------	--------------------	--------------	---------------	----------------	----------------	----------------	----------------	----------------

In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)

All races 3/, both sexes	4.9	1.6	0.3	0.2	0.2	0.4	0.8	1.9	5.1
Male	4.9	1.6	0.4	0.2	0.3	0.4	0.8	2.1	5.7
Female	5.0	1.5	0.3	0.3	0.2	0.4	0.8	1.6	4.5
White, both sexes	5.4	1.6	0.4	0.2	0.2	0.4	0.8	1.8	5.1
Male	5.3	1.6	0.4	0.2	0.3	0.5	0.8	2.1	5.7
Female	5.5	1.6	*	0.2	0.2	0.4	0.8	1.5	4.6
Black, both sexes	3.2	*	*	*	*	0.6	1.1	2.8	5.6
Male	3.1	*	*	*	*	*	1.2	2.9	7.3
Female	3.3	*	*	*	*	0.8	1.0	2.6	4.3

Anemias (D50-D64)

All races 3/, both sexes	1.6	*	0.2	0.1	0.2	0.4	0.4	0.6	1.1
Male	1.2	*	*	0.1	0.2	0.4	0.3	0.6	1.1
Female	2.0	*	*	0.1	0.2	0.4	0.5	0.6	1.0
White, both sexes	1.5	*	*	0.1	0.1	0.1	0.1	0.3	0.8
Male	1.1	*	*	*	*	*	0.1	0.3	0.9
Female	1.9	*	*	*	*	*	0.1	0.3	0.8
Black, both sexes	2.7	*	*	0.5	1.1	2.4	2.5	3.1	3.4
Male	2.4	*	*	*	1.0	2.2	2.1	3.4	3.5
Female	3.1	*	*	*	1.1	2.5	2.8	2.8	3.4

UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

OF DISEASES, 1992), RACE, AND SEX	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS
Malnutrition (E40-E46)									
All races 3/, both sexes	1.4	*	*	*	*	*	0.1	0.2	0.7
Male	1.0	*	*	*	*	*	*	0.2	0.8
Female	1.9	*	*	*	*	*	0.2	0.3	0.6
White, both sexes	1.5	*	*	*	*	*	0.1	0.2	0.6
Male	1.0	*	*	*	*	*	*	0.2	0.6
Female	2.0	*	*	*	*	*	0.1	0.2	0.6
Black, both sexes	1.4	*	*	*	*	*	*	0.6	1.5
Male	1.2	*	*	*	*	*	*	*	2.5
Female	1.6	*	*	*	*	*	*	*	*

Other nutritional deficiencies (E50-E64)

All races 3/, both sexes	0.1	*	*	*	*	*	*	*	*
Male	0.1	*	*	*	*	*	*	*	*
Female	0.1	*	*	*	*	*	*	*	*
White, both sexes	0.1	*	*	*	*	*	*	*	*
Male	0.1	*	*	*	*	*	*	*	*
Female	0.2	*	*	*	*	*	*	*	*
Black, both sexes	0.1	*	*	*	*	*	*	*	*
Male	*	*	*	*	*	*	*	*	*
Female	0.1	*	*	*	*	*	*	*	*

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS
----------------	--------------------	--------------	---------------	----------------	----------------	----------------	----------------	----------------

Alzheimer's disease (G30)

All races 3/, both sexes	18.0	*	*	*	*	*	*	0.2	2.0
Male	10.7	*	*	*	*	*	*	0.2	1.8
Female	25.0	*	*	*	*	*	*	0.2	2.2
White, both sexes	20.5	*	*	*	*	*	*	0.2	2.2
Male	12.2	*	*	*	*	*	*	0.2	1.9
Female	28.6	*	*	*	*	*	*	0.2	2.4
Black, both sexes	7.7	*	*	*	*	*	*	*	1.2
Male	4.6	*	*	*	*	*	*	*	*
Female	10.5	*	*	*	*	*	*	*	*

Major cardiovascular diseases (I00-I78)

All races 3/, both sexes	340.4	16.5	1.5	0.9	3.4	10.1	37.0	117.3	323.2
Male	323.8	17.4	1.7	1.0	4.0	12.8	49.8	168.0	449.7
Female	356.2	15.7	1.4	0.8	2.8	7.5	24.4	68.7	207.8
White, both sexes	360.8	14.3	1.3	0.8	2.8	8.4	31.7	102.9	295.4
Male	340.9	15.0	1.6	0.9	3.4	11.0	44.2	151.9	416.9
Female	379.9	13.6	1.1	0.6	2.2	5.8	19.0	54.9	182.0
Black, both sexes	295.4	29.4	2.8	1.5	6.8	22.4	78.1	247.1	608.0
Male	286.0	30.7	2.4	1.5	7.5	26.5	96.5	330.8	832.4
Female	303.9	28.0	3.3	1.6	6.2	18.7	61.7	177.3	436.9

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS
----------------	--------------------	--------------	---------------	----------------	----------------	----------------	----------------	----------------

Diseases of heart (I00-I09,I11,I13,I20-I51)

All races 3/, both sexes	258.2	12.8	1.2	0.7	2.7	7.9	29.4	95.5	264.5
Male	256.1	13.2	1.4	0.8	3.2	10.3	41.6	142.7	378.6
Female	260.2	12.4	1.0	0.6	2.1	5.5	17.2	50.3	160.4
White, both sexes	274.8	11.3	1.0	0.6	2.2	6.5	25.8	86.2	245.4
Male	271.2	11.7	1.3	0.7	2.6	8.8	37.9	132.2	356.2
Female	278.3	10.9	0.8	0.4	1.6	4.2	13.7	41.2	142.2
Black, both sexes	219.6	20.9	2.2	1.2	5.7	17.9	57.9	186.0	472.0
Male	219.0	20.9	2.1	1.3	6.6	22.0	74.9	257.5	659.4
Female	220.1	21.0	2.3	1.1	4.8	14.2	42.9	126.3	329.1

Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)

All races 3/, both sexes	1.3	*	*	*	*	0.1	0.3	0.7	1.8
Male	0.8	*	*	*	*	*	0.3	0.6	1.5
Female	1.8	*	*	*	*	0.1	0.4	0.8	2.0
White, both sexes	1.4	*	*	*	*	0.1	0.3	0.6	1.7
Male	0.9	*	*	*	*	*	0.2	0.6	1.5
Female	2.0	*	*	*	*	*	0.3	0.7	1.9
Black, both sexes	0.8	*	*	*	*	*	0.6	1.0	2.1
Male	0.5	*	*	*	*	*	*	*	*
Female	1.0	*	*	*	*	*	0.8	1.3	2.6

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS
----------------	--------------------	--------------	---------------	----------------	----------------	----------------	----------------	----------------

Hypertensive heart disease (I11)

All races 3/, both sexes	8.6	*	*	*	0.1	0.6	2.2	5.7	11.1
Male	7.3	*	*	*	0.1	0.7	2.9	7.9	14.7
Female	9.9	*	*	*	*	0.4	1.5	3.5	7.9
White, both sexes	7.7	*	*	*	*	0.3	1.5	3.7	7.7
Male	6.2	*	*	*	*	0.5	2.1	5.6	10.5
Female	9.1	*	*	*	*	0.2	0.9	1.9	5.1
Black, both sexes	16.7	*	*	*	*	2.2	7.4	21.7	42.7
Male	16.2	*	*	*	*	2.7	9.2	28.9	57.4
Female	17.1	*	*	*	*	1.7	5.8	15.6	31.6

Hypertensive heart and renal disease (I13)

All races 3/, both sexes	1.0	*	*	*	*	0.1	0.2	0.4	0.9
Male	0.9	*	*	*	*	*	0.2	0.5	0.9
Female	1.2	*	*	*	*	*	0.2	0.3	0.8
White, both sexes	0.8	*	*	*	*	*	0.1	0.2	0.4
Male	0.7	*	*	*	*	*	0.1	0.2	0.5
Female	1.0	*	*	*	*	*	*	0.1	0.4
Black, both sexes	2.3	*	*	*	*	*	0.8	2.2	4.7
Male	2.1	*	*	*	*	*	1.0	2.9	5.6
Female	2.4	*	*	*	*	*	*	1.6	4.0

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS
----------------	--------------------	--------------	---------------	----------------	----------------	----------------	----------------	----------------

Ischemic heart diseases (I20-I25)

All races 3/, both sexes	187.2	0.8	*	*	0.3	2.5	16.7	67.0	199.1
Male	193.6	*	*	*	0.4	3.8	25.9	105.4	295.6
Female	181.1	*	*	*	0.2	1.2	7.7	30.2	111.2
White, both sexes	201.3	0.8	*	*	0.3	2.3	15.9	63.4	189.0
Male	207.7	*	*	*	0.4	3.5	25.2	101.7	283.8
Female	195.1	*	*	*	0.2	1.0	6.5	25.8	100.7
Black, both sexes	145.2	*	*	*	0.4	4.4	25.8	108.2	318.6
Male	146.8	*	*	*	*	6.1	35.9	155.7	458.8
Female	143.8	*	*	*	*	2.9	16.9	68.6	211.7

Acute myocardial infarction (I21-I22)

All races 3/, both sexes	70.1	0.5	*	*	0.1	1.0	7.0	29.3	87.7
Male	74.5	*	*	*	0.2	1.4	10.7	45.5	128.0
Female	65.8	*	*	*	0.1	0.6	3.3	13.8	50.8
White, both sexes	75.4	*	*	*	0.1	0.9	6.8	28.4	85.7
Male	80.4	*	*	*	0.2	1.4	10.6	45.2	126.6
Female	70.6	*	*	*	*	0.5	2.9	12.0	47.6
Black, both sexes	54.1	*	*	*	*	1.8	10.0	42.1	121.7
Male	53.9	*	*	*	*	2.3	13.5	57.2	167.5
Female	54.3	*	*	*	*	1.4	6.8	29.4	86.7

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS
----------------	--------------------	--------------	---------------	----------------	----------------	----------------	----------------	----------------

Other acute ischemic heart diseases (I24)

All races 3/, both sexes	1.2	*	*	*	*	*	0.2	0.8	2.0
Male	1.3	*	*	*	*	*	0.3	1.2	3.0
Female	1.1	*	*	*	*	*	0.1	0.3	1.1
White, both sexes	1.3	*	*	*	*	*	0.2	0.7	1.8
Male	1.3	*	*	*	*	*	0.3	1.1	2.7
Female	1.2	*	*	*	*	*	*	0.3	0.9
Black, both sexes	1.3	*	*	*	*	*	0.4	1.5	4.3
Male	1.5	*	*	*	*	*	*	2.4	6.4
Female	1.1	*	*	*	*	*	*	*	2.6

Other forms of chronic ischemic heart disease (I20,I25)

All races 3/, both sexes	115.9	*	*	*	0.2	1.5	9.6	37.0	109.5
Male	117.7	*	*	*	0.2	2.3	14.9	58.7	164.6
Female	114.1	*	*	*	*	0.6	4.3	16.1	59.2
White, both sexes	124.6	*	*	*	0.2	1.3	8.9	34.2	101.6
Male	126.0	*	*	*	0.2	2.1	14.3	55.4	154.6
Female	123.3	*	*	*	*	0.5	3.5	13.5	52.1
Black, both sexes	89.8	*	*	*	*	2.6	15.5	64.6	192.7
Male	91.3	*	*	*	*	3.7	21.8	96.0	284.9
Female	88.4	*	*	*	*	1.5	9.8	38.5	122.3

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS
----------------	--------------------	--------------	---------------	----------------	----------------	----------------	----------------	----------------

Atherosclerotic cardiovascular disease, so described (I25.0)

All races 3/, both sexes	25.3	*	*	*	*	0.4	3.1	13.4	34.4
Male	26.7	*	*	*	*	0.6	4.9	21.9	53.6
Female	23.9	*	*	*	*	0.2	1.2	5.3	16.9
White, both sexes	25.5	*	*	*	*	0.3	2.8	11.9	29.7
Male	26.5	*	*	*	*	0.6	4.6	19.8	46.9
Female	24.6	*	*	*	*	*	1.0	4.2	13.6
Black, both sexes	29.5	*	*	*	*	0.7	5.2	27.6	79.2
Male	33.5	*	*	*	*	1.0	7.8	43.7	125.3
Female	25.9	*	*	*	*	*	2.8	14.2	44.1

All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)

All races 3/, both sexes	90.6	*	*	*	0.1	1.1	6.5	23.6	75.1
Male	91.1	*	*	*	0.2	1.7	10.0	36.8	111.0
Female	90.1	*	*	*	*	0.5	3.1	10.9	42.3
White, both sexes	99.1	*	*	*	0.1	1.0	6.1	22.3	71.9
Male	99.5	*	*	*	0.1	1.5	9.7	35.6	107.7
Female	98.7	*	*	*	*	0.4	2.5	9.3	38.5
Black, both sexes	60.3	*	*	*	*	1.9	10.3	37.0	113.4
Male	57.9	*	*	*	*	2.7	14.0	52.3	159.5
Female	62.5	*	*	*	*	1.2	7.0	24.3	78.3

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS
----------------	--------------------	--------------	---------------	----------------	----------------	----------------	----------------	----------------

Other heart diseases (I26-I51)

All races 3/, both sexes	60.1	11.9	1.1	0.6	2.2	4.7	9.9	21.8	51.6
Male	53.6	12.1	1.3	0.7	2.6	5.6	12.3	28.3	65.9
Female	66.3	11.6	0.9	0.5	1.8	3.7	7.6	15.5	38.5
White, both sexes	63.5	10.4	0.9	0.5	1.8	3.8	8.1	18.3	46.6
Male	55.7	10.6	1.2	0.6	2.1	4.7	10.3	24.2	59.9
Female	71.1	10.2	0.7	0.4	1.4	2.9	6.0	12.6	34.1
Black, both sexes	54.7	20.1	2.2	1.2	4.8	10.8	23.3	52.8	103.9
Male	53.4	19.9	2.1	1.3	5.4	12.7	28.3	69.3	136.3
Female	55.8	20.3	2.2	1.1	4.2	9.0	18.8	39.1	79.1

Acute and subacute endocarditis (I33)

All races 3/, both sexes	0.4	*	*	*	*	0.1	0.3	0.5	0.7
Male	0.5	*	*	*	*	0.1	0.3	0.7	0.8
Female	0.4	*	*	*	*	*	0.2	0.3	0.6
White, both sexes	0.4	*	*	*	*	0.1	0.2	0.4	0.6
Male	0.5	*	*	*	*	*	0.3	0.5	0.8
Female	0.3	*	*	*	*	*	0.2	0.2	0.4
Black, both sexes	0.6	*	*	*	*	*	0.5	1.4	1.6
Male	0.7	*	*	*	*	*	0.8	1.9	*
Female	0.6	*	*	*	*	*	*	1.0	1.5

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS
----------------	--------------------	--------------	---------------	----------------	----------------	----------------	----------------	----------------

Diseases of pericardium and acute myocarditis (I30-I31,I40)

All races 3/, both sexes	0.3	0.5	0.2	0.1	0.1	0.1	0.2	0.2	0.4
Male	0.3	*	*	*	0.1	0.2	0.3	0.3	0.5
Female	0.3	*	*	*	*	0.1	0.1	0.2	0.3
White, both sexes	0.3	*	*	*	0.1	0.1	0.2	0.2	0.3
Male	0.3	*	*	*	*	*	0.2	0.2	0.5
Female	0.2	*	*	*	*	*	*	0.2	0.3
Black, both sexes	0.4	*	*	*	*	*	0.4	0.5	*
Male	0.4	*	*	*	*	*	*	*	*
Female	0.4	*	*	*	*	*	*	*	*

Heart failure (I50)

All races 3/, both sexes	20.2	0.8	*	0.1	0.1	0.2	0.6	2.2	9.2
Male	15.7	1.1	*	*	0.1	0.2	0.7	2.7	11.1
Female	24.6	*	*	*	*	0.1	0.5	1.7	7.6
White, both sexes	22.2	*	*	*	*	0.1	0.4	1.8	8.3
Male	17.0	*	*	*	*	0.1	0.6	2.3	10.1
Female	27.1	*	*	*	*	*	0.3	1.3	6.6
Black, both sexes	14.1	*	*	*	*	0.5	1.6	5.8	18.9
Male	11.6	*	*	*	*	*	1.8	7.3	22.3
Female	16.3	*	*	*	*	*	1.6	4.6	16.2

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS
----------------	--------------------	--------------	---------------	----------------	----------------	----------------	----------------	----------------

All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)

All races 3/, both sexes	39.2	10.5	0.9	0.5	2.0	4.3	8.9	18.9	41.3
Male	37.0	10.4	1.0	0.6	2.4	5.2	11.0	24.7	53.5
Female	41.2	10.6	0.8	0.4	1.6	3.4	6.8	13.3	30.1
White, both sexes	40.7	9.4	0.8	0.4	1.6	3.5	7.3	16.0	37.3
Male	37.9	9.2	0.9	0.6	2.0	4.3	9.2	21.2	48.6
Female	43.4	9.5	0.6	0.3	1.2	2.6	5.4	10.9	26.8
Black, both sexes	39.6	16.5	1.6	1.0	4.3	9.8	20.6	45.1	82.8
Male	40.7	15.5	1.8	1.1	4.9	11.6	25.2	59.6	111.7
Female	38.6	17.5	*	0.8	3.8	8.1	16.6	33.0	60.7

Essential (primary) hypertension and hypertensive renal disease (I10,I12)

All races 3/, both sexes	6.6	*	*	*	*	0.2	0.8	2.4	6.0
Male	5.1	*	*	*	*	0.2	1.0	3.1	7.2
Female	7.9	*	*	*	*	0.2	0.6	1.7	5.0
White, both sexes	6.1	*	*	*	*	0.1	0.4	1.4	4.1
Male	4.6	*	*	*	*	0.1	0.6	1.9	5.1
Female	7.5	*	*	*	*	*	0.3	0.9	3.3
Black, both sexes	11.0	*	*	*	*	0.7	3.1	9.9	23.2
Male	9.5	*	*	*	*	*	3.7	12.4	29.0
Female	12.4	*	*	*	*	*	2.6	7.8	18.7

UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

[illegible]

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS
----------------	--------------------	--------------	---------------	----------------	----------------	----------------	----------------	----------------

Other diseases of circulatory system (I71-I78)

All races 3/, both sexes	9.5	*	*	*	0.2	0.4	1.0	2.8	9.3
Male	10.5	*	*	*	0.2	0.6	1.2	3.8	13.4
Female	8.5	*	*	*	0.1	0.3	0.8	1.8	5.6
White, both sexes	10.1	*	*	*	0.1	0.4	0.8	2.4	8.9
Male	11.3	*	*	*	0.2	0.5	1.0	3.4	13.0
Female	9.0	*	*	*	*	0.2	0.6	1.4	5.1
Black, both sexes	7.4	*	*	*	*	0.7	2.2	5.7	14.6
Male	7.2	*	*	*	*	0.9	2.9	7.8	19.3
Female	7.5	*	*	*	*	*	1.6	3.9	11.0

Aortic aneurysm and dissection (I71)

All races 3/, both sexes	5.7	*	*	*	0.1	0.3	0.7	1.8	6.2
Male	7.2	*	*	*	0.1	0.4	1.0	2.7	9.7
Female	4.4	*	*	*	*	0.2	0.4	0.9	3.0
White, both sexes	6.3	*	*	*	0.1	0.3	0.6	1.6	6.1
Male	7.9	*	*	*	*	0.4	0.8	2.5	9.6
Female	4.7	*	*	*	*	0.1	0.3	0.7	2.8
Black, both sexes	3.6	*	*	*	*	0.5	1.6	3.3	8.3
Male	4.0	*	*	*	*	*	2.4	4.8	12.0
Female	3.3	*	*	*	*	*	0.9	2.0	5.5

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS
----------------	--------------------	--------------	---------------	----------------	----------------	----------------	----------------	----------------

Other diseases of arteries, arterioles and capillaries (I72-I78)

All races 3/, both sexes	3.7	*	*	*	0.1	0.1	0.3	1.0	3.1
Male	3.3	*	*	*	*	0.1	0.3	1.1	3.7
Female	4.1	*	*	*	*	0.1	0.4	0.9	2.6
White, both sexes	3.9	*	*	*	*	0.1	0.3	0.8	2.8
Male	3.4	*	*	*	*	0.1	0.2	0.9	3.4
Female	4.3	*	*	*	*	*	0.3	0.7	2.3
Black, both sexes	3.7	*	*	*	*	*	0.6	2.3	6.3
Male	3.2	*	*	*	*	*	*	2.9	7.3
Female	4.2	*	*	*	*	*	0.7	1.9	5.5

Other disorders of circulatory system (I80-I99)

All races 3/, both sexes	1.7	0.7	*	*	0.1	0.4	0.7	1.4	2.2
Male	1.5	*	*	*	*	0.3	0.9	1.6	2.3
Female	1.9	*	*	*	0.1	0.4	0.6	1.2	2.0
White, both sexes	1.7	0.7	*	*	0.1	0.3	0.6	1.3	1.9
Male	1.5	*	*	*	*	0.2	0.8	1.5	2.0
Female	1.9	*	*	*	*	0.3	0.5	1.0	1.8
Black, both sexes	2.2	*	*	*	*	0.9	1.5	2.9	4.5
Male	2.0	*	*	*	*	0.9	1.8	3.5	5.4
Female	2.3	*	*	*	*	1.0	1.3	2.3	3.9

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS
----------------	--------------------	--------------	---------------	----------------	----------------	----------------	----------------	----------------

Influenza and pneumonia (J10-J18)

All races 3/, both sexes	23.7	7.5	0.7	0.2	0.5	1.0	2.4	4.8	12.0
Male	21.3	8.3	0.7	0.2	0.5	1.1	2.8	5.9	14.7
Female	26.1	6.7	0.7	0.2	0.5	0.8	1.9	3.7	9.6
White, both sexes	25.6	5.7	0.6	0.2	0.5	0.9	2.1	4.1	11.0
Male	22.5	6.3	0.5	0.2	0.5	0.9	2.5	4.9	13.1
Female	28.6	5.0	0.7	0.2	0.4	0.8	1.7	3.4	9.0
Black, both sexes	17.0	17.7	1.3	*	0.8	1.8	4.5	10.3	22.3
Male	17.4	18.9	1.9	*	0.7	2.4	5.7	14.5	31.3
Female	16.6	16.4	*	*	0.8	1.2	3.5	6.8	15.4

Influenza (J10-J11)

All races 3/, both sexes	0.6	*	*	*	*	*	0.0	0.1	0.4
Male	0.5	*	*	*	*	*	*	0.1	0.4
Female	0.8	*	*	*	*	*	*	0.1	0.4
White, both sexes	0.7	*	*	*	*	*	*	0.1	0.4
Male	0.6	*	*	*	*	*	*	*	0.4
Female	0.9	*	*	*	*	*	*	0.1	0.4
Black, both sexes	0.2	*	*	*	*	*	*	*	*
Male	0.2	*	*	*	*	*	*	*	*
Female	0.2	*	*	*	*	*	*	*	*

UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

OF DISEASES, 1992), RACE, AND SEX	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS
Chronic lower respiratory diseases (J40-J47)									
All races 3/, both sexes	44.3	0.9	0.3	0.4	0.5	0.8	2.1	8.8	44.8
Male	44.6	1.2	0.4	0.4	0.6	0.9	2.0	9.1	48.7
Female	44.1	*	0.3	0.3	0.4	0.7	2.2	8.4	41.3
White, both sexes	49.9	0.8	0.2	0.2	0.3	0.6	1.7	8.3	46.3
Male	49.3	*	*	0.2	0.4	0.6	1.6	8.5	49.2
Female	50.4	*	*	0.2	0.3	0.6	1.8	8.0	43.5
Black, both sexes	21.5	*	0.9	1.1	1.4	2.2	4.8	14.5	44.5
Male	25.3	*	*	1.4	1.7	2.9	4.8	15.7	57.0
Female	18.2	*	*	0.9	1.0	1.6	4.8	13.5	34.9

Bronchitis, chronic and unspecified (J40-J42)

All races 3/, both sexes	0.4	*	*	*	*	*	0.0	0.1	0.3
Male	0.4	*	*	*	*	*	*	*	0.3
Female	0.5	*	*	*	*	*	*	0.1	0.3
White, both sexes	0.5	*	*	*	*	*	*	0.1	0.3
Male	0.4	*	*	*	*	*	*	*	0.3
Female	0.5	*	*	*	*	*	*	*	0.3
Black, both sexes	0.2	*	*	*	*	*	*	*	*
Male	0.2	*	*	*	*	*	*	*	*
Female	0.2	*	*	*	*	*	*	*	*

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS
----------------	--------------------	--------------	---------------	----------------	----------------	----------------	----------------	----------------

Emphysema (J43)

All races 3/, both sexes	6.1	*	*	*	*	*	0.2	1.3	7.5
Male	6.4	*	*	*	*	*	0.2	1.5	8.2
Female	5.7	*	*	*	*	*	0.2	1.0	6.9
White, both sexes	6.9	*	*	*	*	*	0.2	1.2	8.0
Male	7.2	*	*	*	*	*	0.2	1.4	8.5
Female	6.6	*	*	*	*	*	0.2	1.0	7.5
Black, both sexes	2.6	*	*	*	*	*	*	2.1	6.1
Male	3.3	*	*	*	*	*	*	3.0	8.4
Female	1.8	*	*	*	*	*	*	1.3	4.3

Asthma (J45-J46)

All races 3/, both sexes	1.6	*	0.2	0.3	0.4	0.7	1.0	1.7	2.3
Male	1.2	*	0.3	0.4	0.5	0.8	0.9	1.3	1.7
Female	2.0	*	*	0.3	0.3	0.6	1.2	2.0	2.7
White, both sexes	1.4	*	*	0.2	0.3	0.4	0.7	1.1	1.7
Male	1.0	*	*	0.2	0.4	0.5	0.6	1.0	1.2
Female	1.8	*	*	0.2	0.2	0.4	0.8	1.3	2.1
Black, both sexes	3.3	*	*	1.1	1.3	2.0	3.1	5.5	7.0
Male	2.9	*	*	1.3	1.5	2.6	2.9	3.9	6.4
Female	3.7	*	*	0.9	1.0	1.5	3.3	6.8	7.4

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS
----------------	--------------------	--------------	---------------	----------------	----------------	----------------	----------------	----------------

Pneumonitis due to solids and liquids (J69)

All races 3/, both sexes	6.0	*	*	*	0.1	0.2	0.4	1.0	2.5
Male	6.2	*	*	*	0.1	0.2	0.5	1.2	3.1
Female	5.9	*	*	*	*	0.1	0.3	0.7	1.9
White, both sexes	6.6	*	*	*	0.1	0.2	0.4	0.9	2.4
Male	6.7	*	*	*	*	0.1	0.5	1.1	2.9
Female	6.4	*	*	*	*	0.2	0.3	0.7	1.8
Black, both sexes	4.3	*	*	*	*	*	0.7	1.9	4.2
Male	4.6	*	*	*	*	*	0.8	2.8	5.3
Female	4.1	*	*	*	*	*	*	1.2	3.4

Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)

All races 3/, both sexes	9.3	7.1	0.6	0.2	0.3	0.7	1.4	3.6	10.1
Male	9.2	7.8	0.6	0.2	0.4	0.7	1.5	3.9	11.6
Female	9.3	6.3	0.6	0.2	0.3	0.7	1.2	3.3	8.7
White, both sexes	10.0	5.5	0.5	0.2	0.3	0.6	1.2	3.2	9.8
Male	9.9	6.4	0.5	0.2	0.3	0.6	1.3	3.5	11.2
Female	10.2	4.5	0.6	0.2	0.3	0.6	1.1	3.0	8.4
Black, both sexes	6.7	17.7	0.9	*	0.7	1.1	2.7	7.0	13.6
Male	6.8	17.2	*	*	0.8	1.2	3.2	8.1	15.9
Female	6.6	18.2	*	*	*	1.1	2.2	6.0	11.8

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS
----------------	--------------------	--------------	---------------	----------------	----------------	----------------	----------------	----------------

Hernia (K40-K46)

All races 3/, both sexes	0.6	1.2	*	*	*	*	0.1	0.2	0.5
Male	0.4	1.4	*	*	*	*	*	0.2	0.4
Female	0.7	*	*	*	*	*	*	0.2	0.5
White, both sexes	0.6	1.2	*	*	*	*	0.1	0.2	0.5
Male	0.5	1.5	*	*	*	*	*	0.2	0.4
Female	0.7	*	*	*	*	*	*	0.2	0.5
Black, both sexes	0.4	*	*	*	*	*	*	*	*
Male	0.3	*	*	*	*	*	*	*	*
Female	0.5	*	*	*	*	*	*	*	*

Chronic liver disease and cirrhosis (K70,K73-K74)

All races 3/, both sexes	9.6	*	*	*	0.1	1.1	7.5	17.9	24.1
Male	12.8	*	*	*	*	1.5	10.4	27.4	35.0
Female	6.6	*	*	*	*	0.7	4.7	8.8	14.1
White, both sexes	10.2	*	*	*	0.1	1.0	7.4	17.6	24.0
Male	13.5	*	*	*	*	1.5	10.4	27.1	34.9
Female	6.9	*	*	*	*	0.6	4.5	8.3	13.8
Black, both sexes	7.8	*	*	*	*	1.1	7.9	22.7	26.2
Male	10.3	*	*	*	*	1.1	10.6	34.0	39.7
Female	5.4	*	*	*	*	1.1	5.5	13.2	16.0

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS
----------------	--------------------	--------------	---------------	----------------	----------------	----------------	----------------	----------------

Alcoholic liver disease (K70)

All races 3/, both sexes	4.4	*	*	*	*	0.7	5.0	10.5	12.4
Male	6.7	*	*	*	*	1.1	6.9	16.3	19.4
Female	2.2	*	*	*	*	0.4	3.1	4.8	6.0
White, both sexes	4.5	*	*	*	*	0.7	4.9	10.4	12.3
Male	7.0	*	*	*	*	1.1	6.8	16.4	19.4
Female	2.2	*	*	*	*	0.3	3.0	4.5	5.7
Black, both sexes	3.9	*	*	*	*	0.6	5.0	12.3	13.5
Male	5.5	*	*	*	*	*	7.0	18.2	20.2
Female	2.5	*	*	*	*	*	3.3	7.4	8.3

Other chronic liver disease and cirrhosis (K73-K74)

All races 3/, both sexes	5.2	*	*	*	0.1	0.4	2.6	7.4	11.7
Male	6.1	*	*	*	*	0.4	3.5	11.0	15.6
Female	4.4	*	*	*	*	0.3	1.6	4.0	8.1
White, both sexes	5.6	*	*	*	*	0.3	2.5	7.2	11.7
Male	6.5	*	*	*	*	0.4	3.5	10.8	15.5
Female	4.8	*	*	*	*	0.3	1.5	3.8	8.2
Black, both sexes	3.8	*	*	*	*	0.4	2.9	10.4	12.8
Male	4.8	*	*	*	*	*	3.6	15.9	19.5
Female	2.9	*	*	*	*	*	2.2	5.8	7.7

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS
----------------	--------------------	--------------	---------------	----------------	----------------	----------------	----------------	----------------

Cholelithiasis and other disorders of gallbladder (K80-K82)

All races 3/, both sexes	1.0	*	*	*	*	*	0.1	0.3	0.8
Male	0.9	*	*	*	*	*	0.1	0.3	1.0
Female	1.1	*	*	*	*	*	0.1	0.3	0.6
White, both sexes	1.1	*	*	*	*	*	0.1	0.2	0.7
Male	1.0	*	*	*	*	*	0.1	0.3	0.9
Female	1.2	*	*	*	*	*	*	0.2	0.6
Black, both sexes	0.8	*	*	*	*	*	*	0.7	1.2
Male	0.6	*	*	*	*	*	*	*	*
Female	1.0	*	*	*	*	*	*	*	*

Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)

All races 3/, both sexes	13.5	4.2	*	0.1	0.2	0.7	1.6	4.4	12.9
Male	13.2	5.2	*	*	0.2	0.7	1.9	5.4	14.4
Female	13.8	3.1	*	*	0.2	0.6	1.4	3.5	11.6
White, both sexes	13.1	2.9	*	0.1	0.2	0.5	1.1	3.0	9.7
Male	12.9	3.7	*	*	0.2	0.5	1.3	3.6	11.1
Female	13.2	2.0	*	*	0.2	0.4	0.9	2.4	8.5
Black, both sexes	19.6	12.2	*	*	0.4	1.9	5.4	16.2	41.1
Male	18.3	14.2	*	*	*	2.0	6.5	21.2	46.9
Female	20.7	10.1	*	*	*	1.8	4.3	12.0	36.7

UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

OF DISEASES, 1992), RACE, AND SEX	ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS
Renal failure (N17-N19)									
All races 3/, both sexes	13.2	4.0	*	*	0.2	0.6	1.6	4.3	12.6
Male	12.9	5.1	*	*	0.2	0.6	1.8	5.2	14.1
Female	13.5	2.8	*	*	0.2	0.6	1.3	3.4	11.3
White, both sexes	12.8	2.6	*	*	0.2	0.4	1.1	2.9	9.5
Male	12.7	3.5	*	*	0.2	0.4	1.2	3.4	10.8
Female	13.0	1.7	*	*	0.2	0.4	0.9	2.3	8.2
Black, both sexes	19.2	11.7	*	*	0.4	1.8	5.1	15.6	40.6
Male	18.0	14.2	*	*	*	1.9	6.3	20.3	46.2
Female	20.3	9.1	*	*	*	1.7	4.1	11.6	36.3

Other disorders of kidney (N25,N27)

All races 3/, both sexes	0.0	*	*	*	*	*	*	*
Male	0.0	*	*	*	*	*	*	*
Female	0.0	*	*	*	*	*	*	*
White, both sexes	0.0	*	*	*	*	*	*	*
Male	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*
Black, both sexes	*	*	*	*	*	*	*	*
Male	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS
----------------	--------------------	--------------	---------------	----------------	----------------	----------------	----------------	----------------

Infections of kidney (N10-N12,N13.6,N15.1)

All races 3/, both sexes	0.3	*	*	*	*	*	0.0	0.1	0.2
Male	0.2	*	*	*	*	*	*	*	*
Female	0.4	*	*	*	*	*	*	0.1	0.4
White, both sexes	0.3	*	*	*	*	*	*	0.1	0.2
Male	0.2	*	*	*	*	*	*	*	*
Female	0.4	*	*	*	*	*	*	*	0.3
Black, both sexes	0.3	*	*	*	*	*	*	*	*
Male	0.2	*	*	*	*	*	*	*	*
Female	0.4	*	*	*	*	*	*	*	*

Hyperplasia of prostate (N40)

All races 3/, both sexes	0.2	*	*	*	*	*	*	*	*
Male	0.3	*	*	*	*	*	*	*	*
White, both sexes	0.2	*	*	*	*	*	*	*	*
Male	0.3	*	*	*	*	*	*	*	*
Black, both sexes	0.1	*	*	*	*	*	*	*	*
Male	0.2	*	*	*	*	*	*	*	*

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS
----------------	--------------------	--------------	---------------	----------------	----------------	----------------	----------------	----------------

Inflammatory diseases of female pelvic organs (N70-N76)

All races 3/, both sexes	0.0	*	*	*	*	*	*	*
Female	0.1	*	*	*	*	*	*	*
White, both sexes	0.0	*	*	*	*	*	*	*
Female	0.1	*	*	*	*	*	*	*
Black, both sexes	0.1	*	*	*	*	*	*	*
Female	0.1	*	*	*	*	*	*	*

Pregnancy, childbirth and the puerperium (000-099)

All races 3/, both sexes	0.1	*	0.2	0.5	0.3	*	...
Female	0.3	*	0.5	1.0	0.5	*	...
White, both sexes	0.1	*	0.2	0.4	0.2	*	...
Female	0.2	*	0.4	0.7	0.4	*	...
Black, both sexes	0.4	*	0.6	1.2	0.7	*	...
Female	0.8	*	1.2	2.4	1.3	*	...

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS
----------------	--------------------	--------------	---------------	----------------	----------------	----------------	----------------	----------------

Pregnancy with abortive outcome (000-007)

All races 3/, both sexes	0.0	*	*	*	*	*	...
Female	0.0	*	*	*	*	*	...
White, both sexes	*	*	*	*	*	*	...
Female	*	*	*	*	*	*	...
Black, both sexes	*	*	*	*	*	*	...
Female	*	*	*	*	*	*	...

Other complications of pregnancy, childbirth and the puerperium (010-099)

All races 3/, both sexes	0.1	*	0.2	0.4	0.2	*	...
Female	0.3	*	0.5	0.9	0.5	*	...
White, both sexes	0.1	*	0.2	0.3	0.2	*	...
Female	0.2	*	0.4	0.7	0.4	*	...
Black, both sexes	0.3	*	0.5	1.1	0.6	*	...
Female	0.7	*	1.1	2.0	1.1	*	...

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS
----------------	--------------------	--------------	---------------	----------------	----------------	----------------	----------------	----------------

Certain conditions originating in the perinatal period (P00-P96)

All races 3/, both sexes	5.1	361.7	0.5	0.1	*	*	*	*
Male	5.9	400.6	0.7	0.1	*	*	*	*
Female	4.3	321.2	0.4	*	*	*	*	*
White, both sexes	3.8	278.1	0.4	0.1	*	*	*	*
Male	4.3	307.0	0.5	0.1	*	*	*	*
Female	3.2	247.7	*	*	*	*	*	*
Black, both sexes	14.2	854.7	1.2	*	*	*	*	*
Male	17.0	950.6	*	*	*	*	*	*
Female	11.8	755.3	*	*	*	*	*	*

Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)

All races 3/, both sexes	3.8	149.3	3.3	1.0	1.1	1.3	1.3	1.6	2.4
Male	4.1	154.1	3.3	1.0	1.4	1.5	1.4	1.9	2.3
Female	3.6	144.2	3.2	1.0	0.9	1.1	1.1	1.4	2.4
White, both sexes	3.7	144.2	2.9	0.9	1.1	1.3	1.2	1.7	2.4
Male	3.9	149.3	2.9	0.9	1.3	1.5	1.4	1.9	2.4
Female	3.5	138.7	2.8	1.0	0.9	1.1	1.1	1.5	2.4
Black, both sexes	5.1	189.2	5.7	1.5	1.6	1.5	1.6	1.6	2.4
Male	5.6	191.3	6.2	1.6	2.0	1.9	1.9	2.2	*
Female	4.6	187.0	5.1	1.3	1.1	1.1	1.4	1.2	3.0

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS
----------------	--------------------	--------------	---------------	----------------	----------------	----------------	----------------	----------------

Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

All races 3/, both sexes	11.6	94.6	1.6	0.4	2.3	4.2	6.7	8.5	9.4
Male	11.8	109.7	1.8	0.5	3.0	5.7	8.9	11.7	12.5
Female	11.4	78.8	1.5	0.4	1.5	2.6	4.4	5.4	6.6
White, both sexes	11.6	75.7	1.5	0.4	2.2	4.0	6.5	7.8	8.7
Male	11.5	87.9	1.7	0.4	2.9	5.5	8.9	10.8	11.3
Female	11.7	62.9	1.3	0.3	1.3	2.4	4.1	4.9	6.2
Black, both sexes	14.2	208.1	2.6	0.7	3.2	6.0	9.3	15.1	17.6
Male	16.1	239.5	2.6	0.9	3.7	8.0	10.9	21.5	26.3
Female	12.5	175.5	2.6	*	2.7	4.3	7.8	9.8	10.9

All other diseases (Residual) (D65-E07, E15-E34, E65-F99, G04-G12, G23-G25, G31-H93, K00-K22, K29-K31, K50-K66, K71-K72, K75-K76, K83-M99, N13.0-N13.5, N13.7-N13.9, N14, N15.0, N15.8-N15.9, N20-N23, N28-N39, N41-N64, N80-N98)

All races 3/, both sexes	64.8	28.7	4.0	2.2	4.6	8.3	18.1	35.9	63.2
Male	53.7	32.0	4.4	2.3	5.4	9.7	21.5	43.5	72.0
Female	75.5	25.3	3.7	2.1	3.8	6.8	14.7	28.5	55.2
White, both sexes	68.1	24.4	3.5	2.1	4.3	7.4	16.0	32.3	59.1
Male	55.2	26.4	3.9	2.3	5.2	9.0	19.5	39.8	66.8
Female	80.6	22.3	3.1	1.9	3.4	5.7	12.5	24.9	52.0
Black, both sexes	60.3	52.4	6.7	3.0	6.6	15.6	34.5	71.0	108.8
Male	56.4	63.1	6.9	2.7	7.0	16.3	39.0	84.4	136.1
Female	63.8	41.2	6.4	3.3	6.2	15.0	30.5	59.7	88.1

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS
----------------	--------------------	--------------	---------------	----------------	----------------	----------------	----------------	----------------

Accidents (unintentional injuries) (V01-X59,Y85-Y86)

All races 3/, both sexes	35.6	22.9	12.1	7.5	36.8	31.4	34.3	33.0	31.3
Male	47.4	25.5	14.2	9.2	53.3	48.7	51.3	49.9	44.9
Female	24.2	20.2	9.8	5.8	19.5	14.4	17.6	16.9	18.9
White, both sexes	36.5	18.8	11.1	7.1	38.8	31.9	34.3	31.6	30.4
Male	48.0	21.3	13.1	8.6	56.0	49.2	51.1	47.3	42.7
Female	25.4	16.1	9.0	5.5	20.6	14.4	17.4	16.3	19.1
Black, both sexes	34.8	47.0	18.2	10.1	29.0	33.8	40.1	48.1	41.3
Male	50.9	50.3	21.1	13.0	43.0	54.0	62.5	78.5	71.9
Female	20.2	43.7	15.2	7.0	14.9	15.6	20.1	22.8	18.1

Transport accidents (V01-V99,Y85)

All races 3/, both sexes	17.0	4.5	4.6	4.8	28.7	19.8	17.1	16.1	16.2
Male	23.9	4.8	5.1	5.6	40.2	29.9	24.9	23.4	22.9
Female	10.4	4.2	4.1	4.0	16.8	9.7	9.3	9.0	10.1
White, both sexes	17.3	4.0	4.3	4.9	30.5	19.7	17.1	15.7	16.1
Male	24.1	4.4	4.7	5.6	42.3	29.6	24.8	22.9	22.3
Female	10.7	3.6	3.8	4.1	17.9	9.7	9.4	8.7	10.4
Black, both sexes	16.5	6.9	6.3	5.1	22.0	22.3	18.7	19.8	17.6
Male	25.0	7.1	7.0	6.2	31.7	35.8	30.0	30.8	30.8
Female	8.9	*	5.7	3.9	12.2	10.1	8.8	10.6	7.5

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS
----------------	--------------------	--------------	---------------	----------------	----------------	----------------	----------------	----------------

Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)

All races 3/, both sexes	0.7	*	*	0.1	0.6	0.8	0.9	1.0	1.1
Male	1.2	*	*	0.2	0.9	1.4	1.6	1.7	1.9
Female	0.2	*	*	*	0.2	0.2	0.2	0.3	0.3
White, both sexes	0.8	*	*	0.2	0.6	0.9	1.0	1.0	1.1
Male	1.3	*	*	0.2	1.0	1.6	1.8	1.8	2.0
Female	0.2	*	*	*	0.3	0.2	0.3	0.3	0.4
Black, both sexes	0.4	*	*	*	*	0.5	0.5	0.6	*
Male	0.7	*	*	*	*	0.9	1.0	1.2	*
Female	*	*	*	*	*	*	*	*	*

Nontransport accidents (W00-X59,Y86)

All races 3/, both sexes	18.6	18.4	7.5	2.7	8.0	11.7	17.3	17.0	15.1
Male	23.5	20.7	9.1	3.6	13.1	18.8	26.4	26.4	22.0
Female	13.9	15.9	5.8	1.7	2.7	4.7	8.3	8.0	8.8
White, both sexes	19.2	14.8	6.8	2.2	8.4	12.2	17.2	15.9	14.3
Male	23.8	17.0	8.4	3.0	13.6	19.5	26.3	24.4	20.4
Female	14.8	12.5	5.2	1.4	2.7	4.7	8.0	7.6	8.7
Black, both sexes	18.2	40.2	11.9	5.0	7.1	11.5	21.3	28.3	23.8
Male	25.8	43.2	14.2	6.8	11.4	18.1	32.6	47.7	41.1
Female	11.4	37.1	9.5	3.2	2.7	5.5	11.4	12.2	10.5

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS
----------------	--------------------	--------------	---------------	----------------	----------------	----------------	----------------	----------------

Accidental drowning and submersion (W65-W74)

All races 3/, both sexes	1.3	1.9	3.3	0.9	1.7	1.1	1.1	1.0	0.9
Male	2.0	2.2	4.1	1.3	3.0	2.0	1.8	1.6	1.6
Female	0.5	1.7	2.4	0.6	0.3	0.3	0.3	0.3	0.3
White, both sexes	1.2	1.6	3.5	0.7	1.5	1.1	1.0	0.9	0.9
Male	1.9	1.8	4.4	0.9	2.6	1.9	1.7	1.5	1.6
Female	0.5	*	2.5	0.4	0.3	0.3	0.3	0.3	0.3
Black, both sexes	1.7	4.5	2.5	2.2	2.3	1.1	1.4	1.3	*
Male	2.8	*	3.1	3.2	4.3	2.1	2.6	2.4	*
Female	0.6	*	1.9	1.1	*	*	*	*	*

Accidental exposure to smoke, fire and flames (X00-X09)

All races 3/, both sexes	1.2	1.0	1.9	0.7	0.5	0.6	0.9	1.2	1.5
Male	1.5	1.0	2.2	0.7	0.6	0.8	1.2	1.7	2.1
Female	1.0	*	1.6	0.6	0.4	0.4	0.6	0.7	1.0
White, both sexes	1.1	0.8	1.3	0.5	0.5	0.6	0.8	1.1	1.3
Male	1.3	*	1.6	0.5	0.6	0.8	1.1	1.5	1.7
Female	0.8	*	1.1	0.5	0.3	0.4	0.5	0.6	0.9
Black, both sexes	2.6	*	5.4	1.6	0.8	1.2	1.7	2.4	4.2
Male	3.3	*	6.3	1.8	0.9	1.6	2.4	3.7	6.6
Female	2.0	*	4.6	1.4	*	0.9	1.1	1.4	2.4

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS
----------------	--------------------	--------------	---------------	----------------	----------------	----------------	----------------	----------------

Accidental poisoning and exposure to noxious substances (X40-X49)

All races 3/, both sexes	4.6	*	0.2	0.1	3.0	6.4	10.4	8.2	2.9
Male	6.8	*	0.3	0.1	4.7	9.8	15.2	12.2	3.5
Female	2.6	*	*	*	1.3	3.0	5.7	4.4	2.3
White, both sexes	4.6	*	*	0.1	3.4	6.8	10.4	7.5	2.6
Male	6.8	*	*	*	5.4	10.5	15.3	11.0	2.9
Female	2.6	*	*	*	1.3	3.1	5.5	4.2	2.4
Black, both sexes	5.7	*	*	*	1.4	5.8	13.1	15.9	5.6
Male	8.5	*	*	*	1.7	8.7	18.9	25.9	10.0
Female	3.2	*	*	*	1.0	3.2	7.9	7.5	2.3

Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)

All races 3/, both sexes	6.3	14.9	1.8	0.7	1.7	2.4	3.2	4.1	5.6
Male	7.4	16.6	2.1	0.9	2.9	4.1	5.4	6.7	8.5
Female	5.3	13.0	1.4	0.4	0.5	0.7	1.1	1.5	3.0
White, both sexes	6.7	11.9	1.6	0.7	1.8	2.5	3.2	3.9	5.4
Male	7.6	13.7	2.0	1.0	3.0	4.3	5.4	6.3	8.0
Female	5.8	10.1	1.2	0.4	0.5	0.7	1.0	1.5	3.0
Black, both sexes	5.6	32.3	2.6	0.8	1.5	2.3	3.5	5.8	8.3
Male	7.6	33.7	2.8	1.1	2.5	3.8	6.0	10.5	14.5
Female	3.9	30.8	2.3	*	*	0.9	1.3	1.9	3.6

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS
----------------	--------------------	--------------	---------------	----------------	----------------	----------------	----------------	----------------

Intentional self-harm (suicide) (X60-X84,Y87.0)

All races 3/, both sexes	10.7	0.8	10.4	12.8	14.6	14.6	12.3
Male	17.5	1.2	17.4	21.2	22.9	22.8	19.8
Female	4.1	0.3	3.0	4.5	6.4	6.8	5.4
White, both sexes	11.7	0.8	10.9	13.7	16.1	16.1	13.5
Male	19.1	1.3	18.2	22.4	25.0	24.8	21.5
Female	4.5	0.4	3.1	4.8	7.2	7.6	6.0
Black, both sexes	5.6	0.6	8.2	9.1	7.7	6.7	4.3
Male	9.8	1.1	14.2	16.2	13.3	11.7	7.9
Female	1.8	*	2.2	2.7	2.7	2.5	1.5

Intentional self-harm (suicide) by discharge of firearms (X72-X74)

All races 3/, both sexes	6.0	0.3	5.9	6.6	7.1	7.7	7.6
Male	10.7	0.4	10.4	11.4	11.9	13.3	13.7
Female	1.5	0.1	1.2	1.8	2.4	2.4	2.1
White, both sexes	6.7	0.3	6.2	7.0	8.0	8.6	8.4
Male	11.9	0.5	10.8	12.1	13.1	14.6	14.9
Female	1.7	*	1.3	1.9	2.7	2.6	2.4
Black, both sexes	3.1	*	5.3	5.2	3.4	3.3	3.0
Male	5.8	*	9.6	9.7	6.3	6.1	5.9
Female	0.7	*	0.9	1.2	0.8	1.1	*

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS
----------------	--------------------	--------------	---------------	----------------	----------------	----------------	----------------	----------------

Intentional self-harm (suicide) by other and unspecified means and their sequelae (X60-X71,X75-X84,Y87.0)

All races 3/, both sexes	4.6	0.5	4.5	6.2	7.5	6.9	4.6
Male	6.8	0.8	7.0	9.8	11.0	9.6	6.1
Female	2.6	0.2	1.8	2.8	4.0	4.4	3.3
White, both sexes	5.0	0.5	4.7	6.6	8.2	7.5	5.1
Male	7.3	0.7	7.4	10.3	11.8	10.2	6.6
Female	2.8	0.2	1.9	2.9	4.5	4.9	3.6
Black, both sexes	2.5	0.6	2.9	3.9	4.3	3.3	1.3
Male	4.0	1.0	4.6	6.5	7.0	5.6	1.9
Female	1.1	*	1.3	1.5	1.9	1.4	*

Assault (homicide) (X85-Y09,Y87.1)

All races 3/, both sexes	6.1	9.1	2.3	0.9	12.9	11.1	7.2	4.7	3.1
Male	9.5	10.3	2.6	1.1	21.4	18.1	10.4	7.1	4.7
Female	2.8	7.8	2.1	0.7	3.9	4.3	4.0	2.5	1.6
White, both sexes	3.7	6.5	1.7	0.8	6.5	6.1	4.7	3.3	2.4
Male	5.3	8.1	1.9	0.9	10.1	9.0	6.5	4.6	3.5
Female	2.1	4.9	1.4	0.6	2.7	3.1	2.8	2.1	1.4
Black, both sexes	22.3	24.0	6.5	2.0	48.3	42.5	24.1	15.2	8.5
Male	38.6	24.6	6.6	2.3	85.3	76.9	38.8	27.0	15.3
Female	7.5	23.4	6.4	1.7	11.0	11.4	11.1	5.3	3.2

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS
----------------	--------------------	--------------	---------------	----------------	----------------	----------------	----------------	----------------

Assault (homicide) by discharge of firearms (X93-X95)

All races 3/, both sexes	3.9	*	0.2	0.5	10.3	8.2	4.2	2.6	1.5
Male	6.7	*	*	0.7	18.1	14.2	6.7	4.0	2.3
Female	1.3	*	*	0.3	2.2	2.3	1.9	1.2	0.8
White, both sexes	2.1	*	*	0.4	4.7	4.0	2.7	1.8	1.3
Male	3.3	*	*	0.5	7.8	6.4	3.9	2.6	1.8
Female	1.0	*	*	0.2	1.4	1.6	1.4	1.0	0.8
Black, both sexes	16.1	*	*	1.1	41.7	34.6	15.1	8.6	3.5
Male	30.3	*	*	1.7	76.5	65.9	26.9	15.8	6.6
Female	3.3	*	*	*	6.6	6.4	4.6	2.6	*

Assault (homicide) by other and unspecified means and their sequelae (X85-X92,X96-Y09,Y87.1)

All races 3/, both sexes	2.2	8.8	2.2	0.5	2.5	2.9	2.9	2.1	1.5
Male	2.8	9.9	2.3	0.4	3.3	3.8	3.7	3.0	2.4
Female	1.5	7.6	2.0	0.5	1.7	2.0	2.1	1.3	0.8
White, both sexes	1.6	6.3	1.5	0.4	1.8	2.1	2.0	1.5	1.2
Male	2.0	7.9	1.7	0.4	2.3	2.7	2.6	2.0	1.7
Female	1.1	4.7	1.3	0.4	1.3	1.5	1.4	1.1	0.6
Black, both sexes	6.1	22.8	6.2	0.9	6.6	7.9	9.0	6.6	4.9
Male	8.3	22.3	6.2	*	8.8	11.0	11.8	11.3	8.7
Female	4.2	23.4	6.1	1.2	4.4	5.0	6.5	2.8	2.0

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS
----------------	--------------------	--------------	---------------	----------------	----------------	----------------	----------------	----------------

Legal intervention (Y35,Y89.0)

All races 3/, both sexes	0.1	*	*	*	0.2	0.3	0.2	0.1	*
Male	0.3	*	*	*	0.4	0.6	0.4	0.3	*
Female	*	*	*	*	*	*	*	*	*
White, both sexes	0.1	*	*	*	0.1	0.2	0.2	0.1	*
Male	0.2	*	*	*	0.3	0.5	0.3	0.3	*
Female	*	*	*	*	*	*	*	*	*
Black, both sexes	0.3	*	*	*	0.5	0.6	0.6	*	*
Male	0.6	*	*	*	1.0	1.3	1.2	*	*
Female	*	*	*	*	*	*	*	*	*

Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)

All races 3/, both sexes	1.4	1.7	0.3	0.2	1.0	1.8	2.9	2.3	1.0
Male	2.0	1.4	0.3	0.2	1.5	2.6	4.0	3.1	1.3
Female	0.8	2.0	*	*	0.5	1.0	1.8	1.4	0.6
White, both sexes	1.3	1.4	0.2	0.1	1.0	1.9	2.8	2.1	0.9
Male	1.9	*	*	0.2	1.5	2.6	3.8	2.9	1.1
Female	0.8	1.6	*	*	0.5	1.1	1.7	1.4	0.6
Black, both sexes	1.9	*	*	0.3	0.9	2.0	4.1	3.8	1.8
Male	2.9	*	*	*	1.6	3.2	6.0	6.1	3.2
Female	1.0	*	*	*	*	0.9	2.4	1.8	*

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS
----------------	--------------------	--------------	---------------	----------------	----------------	----------------	----------------	----------------

Discharge of firearms, undetermined intent (Y22-Y24)

All races 3/, both sexes	0.1	*	*	*	0.2	0.1	0.1	0.1	*
Male	0.1	*	*	*	0.3	0.2	0.1	*	*
Female	0.0	*	*	*	*	*	*	*	*
White, both sexes	0.1	*	*	*	0.2	0.1	0.1	0.1	*
Male	0.1	*	*	*	0.3	0.2	0.1	*	*
Female	0.0	*	*	*	*	*	*	*	*
Black, both sexes	0.2	*	*	*	0.3	*	*	*	*
Male	0.3	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*	*

Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)

All races 3/, both sexes	1.3	1.7	0.3	0.1	0.8	1.7	2.8	2.2	0.9
Male	1.8	1.4	0.3	0.2	1.1	2.4	3.8	3.0	1.2
Female	0.8	2.0	*	*	0.4	1.0	1.7	1.4	0.6
White, both sexes	1.3	1.4	0.2	0.1	0.8	1.7	2.7	2.1	0.8
Male	1.7	*	*	0.2	1.2	2.4	3.7	2.8	1.1
Female	0.8	1.6	*	*	0.5	1.0	1.7	1.4	0.6
Black, both sexes	1.7	*	*	*	0.5	1.7	4.0	3.7	1.7
Male	2.6	*	*	*	1.0	2.7	5.9	6.0	2.9
Female	0.9	*	*	*	*	0.8	2.3	1.8	*

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

ALL AGES 1/	UNDER 1 YEAR 2/	1-4 YEARS	5-14 YEARS	15-24 YEARS	25-34 YEARS	35-44 YEARS	45-54 YEARS	55-64 YEARS
----------------	--------------------	--------------	---------------	----------------	----------------	----------------	----------------	----------------

Operations of war and their sequelae (Y36,Y89.1)

All races 3/, both sexes	*	*	*	*	*	*	*	*
Male	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*
White, both sexes	*	*	*	*	*	*	*	*
Male	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*
Black, both sexes	*	*	*	*	*	*	*	*
Male	*	*	*	*	*	*	*	*
Female	*	*	*	*	*	*	*	*

Complications of medical and surgical care (Y40-Y84,Y88)

All races 3/, both sexes	1.1	*	0.1	0.1	0.1	0.2	0.3	0.8	1.6
Male	1.1	*	*	*	*	0.3	0.3	0.8	1.6
Female	1.2	*	*	*	*	0.2	0.3	0.8	1.6
White, both sexes	1.1	*	*	0.1	0.1	0.2	0.3	0.7	1.4
Male	1.1	*	*	*	*	0.2	0.3	0.7	1.4
Female	1.1	*	*	*	*	0.2	0.3	0.7	1.3
Black, both sexes	1.4	*	*	*	*	0.4	0.5	1.4	3.6
Male	1.2	*	*	*	*	*	*	1.3	3.3
Female	1.6	*	*	*	*	*	*	1.5	3.8

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

65-74 YEARS	75-84 YEARS	85 YEARS AND OVER
----------------	----------------	----------------------

All Causes

All races 3/, both sexes	2,428.6	5,688.4	15,321.5
Male	3,015.7	6,854.7	16,605.4
Female	1,945.1	4,900.9	14,768.6
White, both sexes	2,375.1	5,652.4	15,532.5
Male	2,949.8	6,817.7	16,897.7
Female	1,893.9	4,860.4	14,948.7
Black, both sexes	3,251.5	6,826.4	14,752.1
Male	4,065.5	8,240.2	15,494.6
Female	2,659.0	5,967.8	14,441.9

Salmonella infections (A01-A02)

All races 3/, both sexes	*	*	*
Male	*	*	*
Female	*	*	*
White, both sexes	*	*	*
Male	*	*	*
Female	*	*	*
Black, both sexes	*	*	*
Male	*	*	*
Female	*	*	*

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

65-74 75-84 85 YEARS
YEARS YEARS AND OVER

Shigellosis and amebiasis (A03,A06)

All races 3/, both sexes	*	*	*
Male	*	*	*
Female	*	*	*
White, both sexes	*	*	*
Male	*	*	*
Female	*	*	*
Black, both sexes	*	*	*
Male	*	*	*
Female	*	*	*

Certain other intestinal infections (A04,A07-A09)

All races 3/, both sexes	1.2	3.8	11.6
Male	1.3	3.9	10.8
Female	1.2	3.6	12.0
White, both sexes	1.3	4.0	12.1
Male	1.3	4.1	11.4
Female	1.2	3.9	12.4
Black, both sexes	*	*	8.2
Male	*	*	*
Female	*	*	*

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

65-74 75-84 85 YEARS
YEARS YEARS AND OVER

Tuberculosis (A16-A19)

All races 3/, both sexes	0.9	2.0	2.7
Male	1.1	2.8	4.3
Female	0.7	1.5	2.0
White, both sexes	0.5	1.4	2.2
Male	0.7	1.7	3.6
Female	0.4	1.2	1.6
Black, both sexes	3.1	6.7	*
Male	3.9	11.8	*
Female	2.4	3.6	*

Respiratory tuberculosis (A16)

All races 3/, both sexes	0.7	1.7	2.3
Male	0.9	2.3	3.9
Female	0.5	1.3	1.6
White, both sexes	0.4	1.2	1.9
Male	0.6	1.5	3.3
Female	0.3	1.0	1.3
Black, both sexes	2.5	5.3	*
Male	3.6	9.4	*
Female	*	*	*

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

65-74 75-84 85 YEARS
YEARS YEARS AND OVER

Other tuberculosis (A17-A19)

All races 3/, both sexes	0.2	0.3	*
Male	*	0.5	*
Female	0.2	0.3	*
White, both sexes	0.1	0.2	*
Male	*	*	*
Female	*	*	*
Black, both sexes	*	*	*
Male	*	*	*
Female	*	*	*

Whooping cough (A37)

All races 3/, both sexes	*	*	*
Male	*	*	*
Female	*	*	*
White, both sexes	*	*	*
Male	*	*	*
Female	*	*	*
Black, both sexes	*	*	*
Male	*	*	*
Female	*	*	*

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

65-74 75-84 85 YEARS
YEARS YEARS AND OVER

Scarlet fever and erysipelas (A38,A46)

All races 3/, both sexes	*	*	*
Male	*	*	*
Female	*	*	*
White, both sexes	*	*	*
Male	*	*	*
Female	*	*	*
Black, both sexes	*	*	*
Male	*	*	*
Female	*	*	*

Meningococcal infection (A39)

All races 3/, both sexes	*	*	*
Male	*	*	*
Female	*	*	*
White, both sexes	*	*	*
Male	*	*	*
Female	*	*	*
Black, both sexes	*	*	*
Male	*	*	*
Female	*	*	*

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

65-74 YEARS	75-84 YEARS	85 YEARS AND OVER
----------------	----------------	----------------------

Septicemia (A40-A41)

All races 3/, both sexes	31.4	80.7	212.9
Male	34.9	90.3	211.6
Female	28.5	74.2	213.4
White, both sexes	27.8	74.0	200.7
Male	31.3	83.2	200.1
Female	24.9	67.8	201.0
Black, both sexes	70.0	176.6	393.6
Male	77.6	201.1	394.8
Female	64.5	161.8	393.1

Syphilis (A50-A53)

All races 3/, both sexes	*	*	*
Male	*	*	*
Female	*	*	*
White, both sexes	*	*	*
Male	*	*	*
Female	*	*	*
Black, both sexes	*	*	*
Male	*	*	*
Female	*	*	*

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

65-74 75-84 85 YEARS
YEARS YEARS AND OVER

Acute poliomyelitis (A80)

All races 3/, both sexes	*	*	*
Male	*	*	*
Female	*	*	*
White, both sexes	*	*	*
Male	*	*	*
Female	*	*	*
Black, both sexes	*	*	*
Male	*	*	*
Female	*	*	*

Arthropod-borne viral encephalitis (A83-A84,A85.2)

All races 3/, both sexes	*	*	*
Male	*	*	*
Female	*	*	*
White, both sexes	*	*	*
Male	*	*	*
Female	*	*	*
Black, both sexes	*	*	*
Male	*	*	*
Female	*	*	*

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

65-74 75-84 85 YEARS
YEARS YEARS AND OVER

Measles (B05)

All races 3/, both sexes	*	*	*
Male	*	*	*
Female	*	*	*
White, both sexes	*	*	*
Male	*	*	*
Female	*	*	*
Black, both sexes	*	*	*
Male	*	*	*
Female	*	*	*

Viral hepatitis (B15-B19)

All races 3/, both sexes	4.7	5.4	3.9
Male	4.8	6.3	5.4
Female	4.5	4.7	3.3
White, both sexes	4.3	5.2	3.8
Male	4.4	6.1	5.4
Female	4.2	4.5	3.2
Black, both sexes	5.5	6.3	*
Male	6.7	7.4	*
Female	4.7	5.7	*

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

65-74 75-84 85 YEARS
YEARS YEARS AND OVER

Human immunodeficiency virus (HIV) disease (B20-B24)

All races 3/, both sexes	2.2	0.7	*
Male	3.8	1.3	*
Female	0.8	0.3	*
White, both sexes	1.0	0.4	*
Male	1.9	0.7	*
Female	0.3	*	*
Black, both sexes	13.2	4.9	*
Male	23.5	9.1	*
Female	5.7	*	*

Malaria (B50-B54)

All races 3/, both sexes	*	*	*
Male	*	*	*
Female	*	*	*
White, both sexes	*	*	*
Male	*	*	*
Female	*	*	*
Black, both sexes	*	*	*
Male	*	*	*
Female	*	*	*

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

65-74 75-84 85 YEARS
YEARS YEARS AND OVER

Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09,B25-B49,B55-B99)

All races 3/, both sexes	5.9	11.2	20.9
Male	7.3	13.4	26.2
Female	4.8	9.7	18.7
White, both sexes	5.8	11.0	20.8
Male	7.1	13.1	25.6
Female	4.7	9.7	18.7
Black, both sexes	7.6	13.6	24.1
Male	9.8	17.6	29.8
Female	6.0	11.1	21.8

Malignant neoplasms (C00-C97)

All races 3/, both sexes	826.4	1,340.8	1,795.6
Male	1,014.0	1,730.9	2,571.9
Female	671.8	1,077.4	1,461.3
White, both sexes	822.6	1,333.8	1,795.6
Male	999.3	1,707.1	2,569.2
Female	674.7	1,080.1	1,464.7
Black, both sexes	980.0	1,595.5	2,004.0
Male	1,303.5	2,283.6	3,012.7
Female	744.5	1,177.6	1,582.6

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

65-74 75-84 85 YEARS
YEARS YEARS AND OVER

Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)

All races 3/, both sexes	10.8	13.2	19.2
Male	16.2	18.9	25.7
Female	6.4	9.3	16.4
White, both sexes	10.4	13.2	19.5
Male	15.2	18.4	26.3
Female	6.3	9.6	16.6
Black, both sexes	15.1	15.3	15.4
Male	27.0	27.9	*
Female	6.4	7.7	14.7

Malignant neoplasm of esophagus (C15)

All races 3/, both sexes	20.2	25.7	25.6
Male	34.5	45.0	45.3
Female	8.4	12.7	17.1
White, both sexes	19.8	25.4	26.3
Male	34.1	44.9	47.1
Female	7.9	12.1	17.4
Black, both sexes	27.5	34.3	22.3
Male	44.9	56.7	35.1
Female	14.8	20.7	16.9

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

65-74 YEARS	75-84 YEARS	85 YEARS AND OVER
----------------	----------------	----------------------

Malignant neoplasm of stomach (C16)

All races 3/, both sexes	16.8	31.1	52.0
Male	24.0	43.2	71.8
Female	10.9	23.0	43.5
White, both sexes	14.7	27.3	46.8
Male	21.2	38.3	65.7
Female	9.3	19.9	38.7
Black, both sexes	31.2	65.1	101.9
Male	44.6	88.5	125.6
Female	21.4	50.9	92.0

Malignant neoplasms of colon, rectum and anus (C18-C21)

All races 3/, both sexes	77.6	146.6	266.8
Male	96.6	176.5	299.5
Female	62.0	126.4	252.8
White, both sexes	76.1	144.4	266.2
Male	94.8	174.0	298.8
Female	60.5	124.2	252.3
Black, both sexes	102.0	193.6	310.8
Male	125.7	234.0	348.0
Female	84.8	169.1	295.3

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

65-74 75-84 85 YEARS
YEARS YEARS AND OVER

Malignant neoplasms of liver and intrahepatic bile ducts (C22)

All races 3/, both sexes	18.9	28.9	32.3
Male	27.4	41.3	42.4
Female	12.0	20.5	27.9
White, both sexes	17.5	27.9	31.6
Male	25.3	40.4	41.1
Female	11.0	19.3	27.6
Black, both sexes	23.0	29.2	31.7
Male	35.7	38.5	44.7
Female	13.8	23.6	26.2

Malignant neoplasm of pancreas (C25)

All races 3/, both sexes	44.6	76.8	95.1
Male	51.1	84.4	97.6
Female	39.3	71.7	94.1
White, both sexes	43.7	75.5	93.6
Male	50.6	83.8	97.6
Female	38.0	69.9	91.9
Black, both sexes	57.6	99.7	120.1
Male	62.9	103.8	101.1
Female	53.7	97.1	128.1

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

65-74 75-84 85 YEARS
YEARS YEARS AND OVER

Malignant neoplasm of larynx (C32)

All races 3/, both sexes	6.7	8.1	6.4
Male	11.5	15.4	16.5
Female	2.8	3.1	2.1
White, both sexes	6.2	7.7	6.5
Male	10.5	14.8	16.8
Female	2.6	2.9	2.1
Black, both sexes	12.4	14.3	6.9
Male	23.2	27.6	*
Female	4.6	6.3	*

Malignant neoplasms of trachea, bronchus and lung (C33-C34)

All races 3/, both sexes	287.6	372.2	298.1
Male	382.5	523.2	494.5
Female	209.5	270.3	213.5
White, both sexes	291.1	375.7	301.1
Male	380.6	521.1	498.3
Female	216.2	276.9	216.8
Black, both sexes	307.9	382.0	288.8
Male	467.2	623.9	519.3
Female	191.9	235.0	192.5

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

65-74 75-84 85 YEARS
YEARS YEARS AND OVER

Malignant melanoma of skin (C43)

All races 3/, both sexes	9.0	13.2	18.7
Male	13.0	20.4	30.2
Female	5.7	8.3	13.8
White, both sexes	10.1	14.3	20.1
Male	14.5	22.2	33.1
Female	6.4	9.0	14.5
Black, both sexes	1.8	3.6	6.6
Male	*	*	*
Female	*	*	*

Malignant neoplasm of breast (C50)

All races 3/, both sexes	49.7	79.2	145.5
Male	1.4	2.2	3.9
Female	89.4	131.2	206.4
White, both sexes	49.9	79.6	145.6
Male	1.4	2.2	3.7
Female	90.5	132.2	206.3
Black, both sexes	57.1	92.2	176.9
Male	*	*	*
Female	97.5	146.4	247.2

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

65-74 YEARS	75-84 YEARS	85 YEARS AND OVER
----------------	----------------	----------------------

Malignant neoplasm of cervix uteri (C53)

All races 3/, both sexes	3.8	4.9	7.2
Female	6.9	8.1	10.3
White, both sexes	3.1	4.3	6.2
Female	5.8	7.2	8.9
Black, both sexes	9.3	12.1	19.8
Female	16.1	19.5	28.0

Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)

All races 3/, both sexes	9.8	15.9	24.4
Female	17.9	26.6	34.9
White, both sexes	8.7	14.9	24.1
Female	15.9	25.0	34.4
Black, both sexes	22.1	31.1	33.2
Female	38.2	50.0	47.1

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

65-74 YEARS	75-84 YEARS	85 YEARS AND OVER
----------------	----------------	----------------------

Malignant neoplasm of ovary (C56)

All races 3/, both sexes	19.8	33.3	40.4
Female	36.1	55.8	57.8
White, both sexes	20.4	34.3	41.5
Female	37.5	57.7	59.3
Black, both sexes	17.6	27.1	34.5
Female	30.4	43.6	48.9

Malignant neoplasm of prostate (C61)

All races 3/, both sexes	37.5	106.2	206.6
Male	83.0	263.4	686.1
White, both sexes	33.0	97.3	196.4
Male	72.3	240.4	655.7
Black, both sexes	87.8	234.1	357.5
Male	208.5	619.5	1,213.2

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

65-74 75-84 85 YEARS
YEARS YEARS AND OVER

Malignant neoplasms of kidney and renal pelvis (C64-C65)

All races 3/, both sexes	17.3	26.7	33.3
Male	24.5	37.6	52.3
Female	11.4	19.3	25.2
White, both sexes	17.7	27.0	34.1
Male	24.8	38.0	53.9
Female	11.8	19.5	25.6
Black, both sexes	16.1	26.1	29.8
Male	25.2	37.0	45.8
Female	9.5	19.5	23.1

Malignant neoplasm of bladder (C67)

All races 3/, both sexes	15.7	35.8	68.7
Male	25.5	61.5	132.2
Female	7.6	18.4	41.4
White, both sexes	16.4	36.5	71.2
Male	26.9	63.7	138.9
Female	7.6	18.1	42.3
Black, both sexes	13.0	35.3	49.9
Male	16.8	51.2	81.9
Female	10.2	25.7	36.5

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

65-74 75-84 85 YEARS
YEARS YEARS AND OVER

Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)

All races 3/, both sexes	16.7	20.1	16.7
Male	20.5	23.9	21.3
Female	13.6	17.6	14.7
White, both sexes	18.0	21.1	17.3
Male	22.0	25.1	22.5
Female	14.7	18.3	15.1
Black, both sexes	7.9	12.1	10.7
Male	9.4	12.6	*
Female	6.8	11.8	11.1

Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)

All races 3/, both sexes	77.3	149.8	201.9
Male	98.0	189.4	267.4
Female	60.2	123.1	173.7
White, both sexes	79.1	153.7	206.3
Male	100.2	194.9	276.8
Female	61.5	125.7	176.2
Black, both sexes	73.1	123.2	175.9
Male	91.7	150.5	207.5
Female	59.5	106.6	162.7

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

65-74 75-84 85 YEARS
YEARS YEARS AND OVER

Hodgkin's disease (C81)

All races 3/, both sexes	1.3	2.1	2.4
Male	1.6	2.4	3.5
Female	1.0	1.8	2.0
White, both sexes	1.3	2.2	2.6
Male	1.7	2.6	3.7
Female	1.0	2.0	2.1
Black, both sexes	*	*	*
Male	*	*	*
Female	*	*	*

Non-Hodgkin's lymphoma (C82-C85)

All races 3/, both sexes	30.8	61.2	79.2
Male	37.8	73.2	100.4
Female	25.0	53.1	70.0
White, both sexes	32.5	64.4	83.4
Male	39.7	77.1	106.7
Female	26.5	55.8	73.4
Black, both sexes	18.9	27.3	36.4
Male	23.8	30.3	40.4
Female	15.3	25.5	34.7

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

65-74 75-84 85 YEARS
YEARS YEARS AND OVER

Leukemia (C91-C95)

All races 3/, both sexes	27.6	54.7	82.5
Male	37.7	74.4	113.7
Female	19.3	41.3	69.0
White, both sexes	28.7	56.7	85.1
Male	39.1	77.2	117.9
Female	20.1	42.8	71.1
Black, both sexes	22.1	38.9	66.2
Male	30.2	53.2	90.5
Female	16.2	30.2	56.0

Multiple myeloma and immunoproliferative neoplasms (C88,C90)

All races 3/, both sexes	17.5	31.6	37.3
Male	20.7	38.9	49.1
Female	14.9	26.7	32.3
White, both sexes	16.4	30.1	34.8
Male	19.5	37.6	47.9
Female	13.8	25.0	29.2
Black, both sexes	31.3	56.1	71.8
Male	36.8	65.9	73.4
Female	27.3	50.2	71.1

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

65-74 75-84 85 YEARS
YEARS YEARS AND OVER

Other and unspecified malignant neoplasms of lymphoid, hematopoietic and related tissue (C96)

All races 3/, both sexes	0.1	0.2	0.5
Male	*	*	*
Female	*	*	*
White, both sexes	0.1	0.2	*
Male	*	*	*
Female	*	*	*
Black, both sexes	*	*	*
Male	*	*	*
Female	*	*	*

All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)

All races 3/, both sexes	86.5	153.2	236.6
Male	104.5	184.6	285.3
Female	71.7	132.0	215.7
White, both sexes	86.5	153.7	241.0
Male	104.7	185.1	292.8
Female	71.3	132.3	218.8
Black, both sexes	97.6	165.0	211.4
Male	117.2	204.3	234.1
Female	83.4	141.1	201.9

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

65-74 YEARS	75-84 YEARS	85 YEARS AND OVER
----------------	----------------	----------------------

In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)

All races 3/, both sexes	14.0	37.7	83.8
Male	17.2	46.7	103.3
Female	11.4	31.6	75.4
White, both sexes	14.1	38.7	86.4
Male	17.2	48.2	107.4
Female	11.5	32.3	77.4
Black, both sexes	14.6	31.6	64.3
Male	18.3	35.0	70.2
Female	11.9	29.5	61.8

Anemias (D50-D64)

All races 3/, both sexes	2.8	9.5	41.8
Male	2.9	9.4	38.0
Female	2.8	9.6	43.4
White, both sexes	2.6	9.3	41.6
Male	2.8	9.1	37.3
Female	2.4	9.4	43.4
Black, both sexes	5.2	13.6	51.1
Male	4.3	13.5	51.1
Female	5.9	13.6	51.1

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

65-74 75-84 85 YEARS
YEARS YEARS AND OVER

Diabetes mellitus (E10-E14)

All races 3/, both sexes	91.8	180.2	315.6
Male	100.3	196.2	327.4
Female	84.8	169.3	310.5
White, both sexes	82.4	166.4	300.1
Male	93.2	187.4	321.8
Female	73.3	152.1	290.8
Black, both sexes	183.7	357.2	526.3
Male	176.8	327.2	417.2
Female	188.7	375.4	571.8

Nutritional deficiencies (E40-E64)

All races 3/, both sexes	2.5	10.4	49.4
Male	2.7	9.5	41.4
Female	2.4	11.1	52.8
White, both sexes	2.3	10.0	48.3
Male	2.4	9.0	40.9
Female	2.3	10.7	51.5
Black, both sexes	4.6	16.4	67.1
Male	6.2	17.3	58.5
Female	3.5	15.9	70.7

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

65-74 75-84 85 YEARS
YEARS YEARS AND OVER

Malnutrition (E40-E46)

All races 3/, both sexes	2.3	9.7	45.7
Male	2.5	8.7	38.6
Female	2.2	10.4	48.8
White, both sexes	2.1	9.4	44.7
Male	2.2	8.3	37.9
Female	2.0	10.1	47.6
Black, both sexes	4.5	15.2	64.0
Male	6.0	16.8	56.4
Female	3.5	14.3	67.1

Other nutritional deficiencies (E50-E64)

All races 3/, both sexes	0.2	0.7	3.6
Male	*	0.7	2.9
Female	0.2	0.7	4.0
White, both sexes	0.2	0.7	3.6
Male	*	0.7	3.0
Female	0.3	0.6	3.9
Black, both sexes	*	*	*
Male	*	*	*
Female	*	*	*

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

65-74 75-84 85 YEARS
YEARS YEARS AND OVER

Meningitis (G00,G03)

All races 3/, both sexes	0.6	1.0	1.2
Male	0.7	1.2	*
Female	0.5	0.8	1.0
White, both sexes	0.5	0.9	1.0
Male	0.7	1.2	*
Female	0.4	0.8	1.0
Black, both sexes	1.6	*	*
Male	*	*	*
Female	*	*	*

Parkinson's disease (G20-G21)

All races 3/, both sexes	11.6	62.1	130.1
Male	16.3	91.3	210.9
Female	7.7	42.5	95.3
White, both sexes	12.5	65.8	138.0
Male	17.4	95.8	224.5
Female	8.3	45.4	101.0
Black, both sexes	5.5	27.9	51.7
Male	8.4	46.5	79.8
Female	3.5	16.6	40.0

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

65-74 YEARS	75-84 YEARS	85 YEARS AND OVER
----------------	----------------	----------------------

Alzheimer's disease (G30)

All races 3/, both sexes	18.9	140.1	659.0
Male	18.9	125.5	495.4
Female	18.8	150.0	729.5
White, both sexes	19.8	145.2	687.6
Male	19.8	130.3	517.1
Female	19.8	155.3	760.5
Black, both sexes	15.4	112.0	446.3
Male	15.8	94.7	342.7
Female	15.1	122.5	489.6

Major cardiovascular diseases (I00-I78)

All races 3/, both sexes	859.7	2,405.5	7,925.3
Male	1,127.6	2,877.1	8,015.6
Female	639.1	2,087.1	7,886.5
White, both sexes	825.9	2,376.9	8,043.5
Male	1,097.1	2,863.9	8,188.3
Female	598.9	2,045.9	7,981.6
Black, both sexes	1,283.0	3,024.3	7,538.2
Male	1,567.4	3,400.3	7,093.9
Female	1,075.9	2,795.8	7,723.8

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

65-74 YEARS	75-84 YEARS	85 YEARS AND OVER
----------------	----------------	----------------------

Diseases of heart (I00-I09,I11,I13,I20-I51)

All races 3/, both sexes	673.8	1,787.1	5,848.7
Male	909.2	2,210.1	6,100.8
Female	479.9	1,501.5	5,740.1
White, both sexes	651.8	1,772.4	5,953.9
Male	891.2	2,209.6	6,257.6
Female	451.3	1,475.2	5,824.0
Black, both sexes	977.3	2,200.0	5,403.4
Male	1,212.8	2,522.4	5,198.6
Female	805.9	2,004.2	5,489.0

Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)

All races 3/, both sexes	4.5	9.6	16.6
Male	2.9	6.5	11.6
Female	5.8	11.6	18.7
White, both sexes	4.6	10.2	17.4
Male	2.9	6.9	12.1
Female	6.1	12.4	19.7
Black, both sexes	3.4	4.0	8.5
Male	*	*	*
Female	4.3	5.0	*

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

65-74 75-84 85 YEARS
YEARS YEARS AND OVER

Hypertensive heart disease (I11)

All races 3/, both sexes	19.8	48.4	190.6
Male	24.0	45.7	132.0
Female	16.4	50.2	215.8
White, both sexes	15.2	41.7	179.8
Male	18.9	38.9	123.4
Female	12.1	43.5	203.9
Black, both sexes	65.2	134.3	340.9
Male	77.8	135.2	252.2
Female	56.0	133.8	378.0

Hypertensive heart and renal disease (I13)

All races 3/, both sexes	2.3	7.0	24.0
Male	2.5	7.1	26.4
Female	2.1	7.0	23.0
White, both sexes	1.5	5.6	21.8
Male	1.7	6.1	23.2
Female	1.4	5.3	21.2
Black, both sexes	9.7	23.2	50.5
Male	10.8	20.3	58.5
Female	8.9	25.0	47.1

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

65-74 YEARS	75-84 YEARS	85 YEARS AND OVER
----------------	----------------	----------------------

Ischemic heart diseases (I20-I25)

All races 3/, both sexes	511.3	1,329.4	4,127.8
Male	713.7	1,707.5	4,461.0
Female	344.6	1,074.0	3,984.3
White, both sexes	500.5	1,324.3	4,210.4
Male	707.8	1,715.7	4,591.4
Female	326.9	1,058.3	4,047.5
Black, both sexes	684.5	1,551.6	3,694.1
Male	867.8	1,822.6	3,585.3
Female	551.1	1,387.0	3,739.6

Acute myocardial infarction (I21-I22)

All races 3/, both sexes	211.7	505.2	1,321.0
Male	290.4	637.3	1,488.0
Female	146.9	416.0	1,249.0
White, both sexes	209.7	503.9	1,338.4
Male	291.5	641.6	1,519.3
Female	141.2	410.4	1,261.1
Black, both sexes	262.0	581.7	1,278.9
Male	321.0	667.4	1,309.0
Female	219.0	529.7	1,266.4

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

65-74 YEARS	75-84 YEARS	85 YEARS AND OVER
----------------	----------------	----------------------

Other acute ischemic heart diseases (I24)

All races 3/, both sexes	3.5	7.4	21.6
Male	4.9	9.0	19.6
Female	2.5	6.3	22.4
White, both sexes	3.3	7.1	21.9
Male	4.5	8.7	20.0
Female	2.4	6.0	22.7
Black, both sexes	5.5	11.0	22.0
Male	8.3	13.5	*
Female	3.5	9.5	23.1

Other forms of chronic ischemic heart disease (I20,I25)

All races 3/, both sexes	296.0	816.8	2,785.2
Male	418.5	1,061.2	2,953.4
Female	195.2	651.8	2,712.8
White, both sexes	287.5	813.3	2,850.1
Male	411.8	1,065.4	3,052.1
Female	183.4	641.9	2,763.7
Black, both sexes	417.0	958.9	2,393.2
Male	538.6	1,141.7	2,257.2
Female	328.6	847.9	2,450.1

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

65-74 YEARS	75-84 YEARS	85 YEARS AND OVER
----------------	----------------	----------------------

Atherosclerotic cardiovascular disease, so described (I25.0)

All races 3/, both sexes	69.2	158.7	527.6
Male	98.4	195.9	522.6
Female	45.1	133.6	529.8
White, both sexes	62.1	149.6	523.6
Male	89.1	184.4	520.4
Female	39.5	125.9	525.0
Black, both sexes	141.9	283.6	648.3
Male	198.2	359.0	647.0
Female	100.9	237.9	648.8

All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)

All races 3/, both sexes	226.9	658.1	2,257.6
Male	320.1	865.3	2,430.8
Female	150.1	518.2	2,183.1
White, both sexes	225.4	663.7	2,326.5
Male	322.7	881.0	2,531.7
Female	143.9	516.0	2,238.8
Black, both sexes	275.2	675.3	1,745.0
Male	340.4	782.7	1,610.1
Female	227.7	610.0	1,801.3

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

65-74 75-84 85 YEARS
YEARS YEARS AND OVER

Other heart diseases (I26-I51)

All races 3/, both sexes	135.8	392.8	1,489.7
Male	166.0	443.4	1,469.8
Female	111.0	358.6	1,498.2
White, both sexes	129.9	390.6	1,524.5
Male	159.8	442.1	1,507.5
Female	104.8	355.6	1,531.7
Black, both sexes	214.6	486.8	1,309.4
Male	254.2	541.9	1,294.1
Female	185.8	453.4	1,315.8

Acute and subacute endocarditis (I33)

All races 3/, both sexes	1.2	2.5	2.8
Male	1.5	3.4	3.9
Female	1.0	1.8	2.3
White, both sexes	1.2	2.4	2.7
Male	1.4	3.4	4.0
Female	1.0	1.8	2.1
Black, both sexes	1.8	2.9	*
Male	*	*	*
Female	*	*	*

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

65-74 YEARS	75-84 YEARS	85 YEARS AND OVER
----------------	----------------	----------------------

Diseases of pericardium and acute myocarditis (I30-I31,I40)

All races 3/, both sexes	0.7	1.2	1.8
Male	0.8	1.5	2.0
Female	0.6	1.0	1.7
White, both sexes	0.7	1.2	1.8
Male	0.7	1.4	2.0
Female	0.6	1.0	1.7
Black, both sexes	*	*	*
Male	*	*	*
Female	*	*	*

Heart failure (I50)

All races 3/, both sexes	35.2	136.6	675.9
Male	41.6	151.1	637.8
Female	30.0	126.8	692.3
White, both sexes	33.9	137.0	695.9
Male	40.7	152.0	658.2
Female	28.3	126.8	712.1
Black, both sexes	54.0	157.3	559.8
Male	59.2	168.5	536.4
Female	50.3	150.5	569.6

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

65-74 YEARS	75-84 YEARS	85 YEARS AND OVER
----------------	----------------	----------------------

All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)

All races 3/, both sexes	98.6	252.6	809.2
Male	122.1	287.4	826.1
Female	79.3	229.0	802.0
White, both sexes	94.1	250.0	824.1
Male	117.0	285.2	843.3
Female	74.9	226.1	815.9
Black, both sexes	157.7	325.5	743.9
Male	191.3	368.7	753.4
Female	133.2	299.3	739.9

Essential (primary) hypertension and hypertensive renal disease (I10,I12)

All races 3/, both sexes	15.3	45.7	160.8
Male	15.9	44.4	135.1
Female	14.8	46.5	171.9
White, both sexes	12.0	40.5	151.5
Male	12.8	39.7	126.4
Female	11.3	41.1	162.3
Black, both sexes	47.5	109.1	284.1
Male	48.2	105.8	249.0
Female	46.9	111.1	298.8

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

65-74 YEARS	75-84 YEARS	85 YEARS AND OVER
----------------	----------------	----------------------

Cerebrovascular diseases (I60-I69)

All races 3/, both sexes	130.2	463.1	1,568.4
Male	146.8	482.5	1,408.3
Female	116.5	450.0	1,637.4
White, both sexes	121.1	453.5	1,583.8
Male	135.9	472.0	1,421.8
Female	108.8	441.0	1,653.1
Black, both sexes	217.3	599.8	1,533.9
Male	256.6	640.1	1,346.2
Female	188.7	575.4	1,612.4

Atherosclerosis (I70)

All races 3/, both sexes	7.8	34.7	186.3
Male	9.8	39.0	167.5
Female	6.3	31.9	194.5
White, both sexes	7.9	35.2	193.0
Male	9.9	39.7	173.0
Female	6.2	32.2	201.6
Black, both sexes	8.7	35.9	138.6
Male	10.5	39.1	136.2
Female	7.4	33.9	139.6

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

65-74 75-84 85 YEARS
YEARS YEARS AND OVER

Other diseases of circulatory system (I71-I78)

All races 3/, both sexes	32.6	74.9	161.1
Male	46.0	101.1	203.9
Female	21.6	57.2	142.6
White, both sexes	33.2	75.3	161.2
Male	47.4	102.9	209.5
Female	21.3	56.5	140.6
Black, both sexes	32.2	79.4	178.1
Male	39.3	92.9	163.9
Female	27.0	71.3	184.1

Aortic aneurysm and dissection (I71)

All races 3/, both sexes	22.1	47.4	78.1
Male	33.3	69.4	120.8
Female	12.9	32.5	59.8
White, both sexes	23.0	48.6	79.9
Male	35.0	71.8	126.6
Female	12.9	32.8	59.9
Black, both sexes	16.6	37.0	62.4
Male	21.1	45.6	63.9
Female	13.2	31.8	61.8

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

65-74 75-84 85 YEARS
YEARS YEARS AND OVER

Other diseases of arteries, arterioles and capillaries (I72-I78)

All races 3/, both sexes	10.5	27.5	82.9
Male	12.7	31.8	83.1
Female	8.8	24.6	82.9
White, both sexes	10.2	26.7	81.4
Male	12.4	31.1	82.9
Female	8.3	23.7	80.7
Black, both sexes	15.6	42.4	115.7
Male	18.2	47.3	100.0
Female	13.7	39.5	122.3

Other disorders of circulatory system (I80-I99)

All races 3/, both sexes	4.4	9.4	25.2
Male	4.8	9.5	21.1
Female	4.1	9.4	27.0
White, both sexes	4.0	9.2	25.2
Male	4.5	9.3	21.0
Female	3.6	9.0	27.0
Black, both sexes	8.7	14.7	30.1
Male	8.7	12.9	27.7
Female	8.7	15.7	31.1

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

65-74 YEARS	75-84 YEARS	85 YEARS AND OVER
----------------	----------------	----------------------

Influenza and pneumonia (J10-J18)

All races 3/, both sexes	39.6	161.0	734.4
Male	48.9	196.5	846.3
Female	31.9	137.0	686.2
White, both sexes	37.6	160.4	749.9
Male	45.8	194.5	864.2
Female	30.7	137.2	701.0
Black, both sexes	58.7	174.3	602.5
Male	79.9	227.3	697.0
Female	43.4	142.1	562.9

Influenza (J10-J11)

All races 3/, both sexes	1.1	4.0	19.9
Male	1.3	3.9	19.6
Female	0.9	4.0	20.0
White, both sexes	1.1	4.2	21.1
Male	1.4	4.2	20.6
Female	0.9	4.3	21.3
Black, both sexes	*	*	9.4
Male	*	*	*
Female	*	*	*

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

65-74 75-84 85 YEARS
YEARS YEARS AND OVER

Pneumonia (J12-J18)

All races 3/, both sexes	38.5	157.0	714.5
Male	47.6	192.5	826.7
Female	30.9	133.0	666.1
White, both sexes	36.5	156.2	728.8
Male	44.4	190.4	843.6
Female	29.8	132.9	679.7
Black, both sexes	57.8	172.9	593.1
Male	79.2	225.8	685.3
Female	42.2	140.7	554.5

Other acute lower respiratory infections (J20-J22)

All races 3/, both sexes	0.2	0.7	4.5
Male	0.3	0.7	3.6
Female	0.2	0.6	4.8
White, both sexes	0.3	0.7	4.7
Male	0.3	0.7	3.6
Female	0.2	0.7	5.1
Black, both sexes	*	*	*
Male	*	*	*
Female	*	*	*

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

65-74 75-84 85 YEARS
YEARS YEARS AND OVER

Acute bronchitis and bronchiolitis (J20-J21)

All races 3/, both sexes	0.2	0.4	2.6
Male	*	0.5	2.2
Female	*	0.4	2.7
White, both sexes	0.2	0.4	2.7
Male	*	0.6	2.3
Female	*	0.3	2.9
Black, both sexes	*	*	*
Male	*	*	*
Female	*	*	*

Unspecified acute lower respiratory infection (J22)

All races 3/, both sexes	*	0.2	1.9
Male	*	*	*
Female	*	0.3	2.1
White, both sexes	*	0.3	2.0
Male	*	*	*
Female	*	0.3	2.2
Black, both sexes	*	*	*
Male	*	*	*
Female	*	*	*

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

65-74 YEARS	75-84 YEARS	85 YEARS AND OVER
----------------	----------------	----------------------

Chronic lower respiratory diseases (J40-J47)

All races 3/, both sexes	171.5	387.6	640.1
Male	197.6	480.3	918.3
Female	150.0	324.9	520.2
White, both sexes	181.2	405.4	668.1
Male	204.4	495.2	951.4
Female	161.8	344.4	547.0
Black, both sexes	118.6	239.1	374.8
Male	169.1	359.6	639.6
Female	81.9	165.9	264.1

Bronchitis, chronic and unspecified (J40-J42)

All races 3/, both sexes	0.9	2.7	11.7
Male	1.0	3.5	12.2
Female	0.8	2.2	11.4
White, both sexes	0.9	2.8	12.5
Male	1.0	3.7	13.2
Female	0.8	2.2	12.3
Black, both sexes	*	2.2	*
Male	*	*	*
Female	*	*	*

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

65-74 75-84 85 YEARS
YEARS YEARS AND OVER

Emphysema (J43)

All races 3/, both sexes	28.2	52.9	62.5
Male	33.4	68.0	98.8
Female	23.9	42.7	46.9
White, both sexes	30.1	55.8	65.1
Male	34.8	70.9	100.7
Female	26.2	45.4	49.8
Black, both sexes	16.6	27.4	37.3
Male	26.2	40.3	80.9
Female	9.6	19.6	19.1

Asthma (J45-J46)

All races 3/, both sexes	4.0	6.9	16.5
Male	3.1	4.9	12.2
Female	4.8	8.2	18.3
White, both sexes	3.4	6.5	15.7
Male	2.6	4.6	10.8
Female	4.2	7.8	17.8
Black, both sexes	9.0	10.6	22.3
Male	7.6	6.2	23.4
Female	10.1	13.2	21.8

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

65-74 75-84 85 YEARS
YEARS YEARS AND OVER

Other chronic lower respiratory diseases (J44,J47)

All races 3/, both sexes	138.4	325.1	549.5
Male	160.2	403.9	795.1
Female	120.5	271.8	443.7
White, both sexes	146.8	340.3	574.8
Male	166.0	416.0	826.7
Female	130.6	288.9	467.1
Black, both sexes	92.1	198.9	311.4
Male	134.5	311.1	532.1
Female	61.3	130.7	219.2

Pneumoconioses and chemical effects (J60-J66,J68)

All races 3/, both sexes	1.5	3.9	7.4
Male	3.2	9.4	23.7
Female	*	*	*
White, both sexes	1.7	4.1	7.8
Male	3.5	9.8	25.2
Female	*	*	*
Black, both sexes	*	2.6	*
Male	*	6.2	*
Female	*	*	*

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

65-74 YEARS	75-84 YEARS	85 YEARS AND OVER
----------------	----------------	----------------------

Pneumonitis due to solids and liquids (J69)

All races 3/, both sexes	10.5	44.7	185.2
Male	14.6	61.2	261.9
Female	7.1	33.6	152.1
White, both sexes	10.0	44.6	188.9
Male	13.6	61.1	270.9
Female	6.9	33.3	153.8
Black, both sexes	17.1	52.4	160.3
Male	26.6	72.3	193.7
Female	10.2	40.4	146.3

Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)

All races 3/, both sexes	30.0	72.3	139.9
Male	36.1	89.0	170.5
Female	25.0	61.0	126.7
White, both sexes	30.5	73.9	144.3
Male	36.6	91.0	175.9
Female	25.3	62.3	130.8
Black, both sexes	29.7	59.1	98.5
Male	34.4	70.9	122.4
Female	26.3	52.0	88.5

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

65-74 75-84 85 YEARS
YEARS YEARS AND OVER

Peptic ulcer (K25-K28)

All races 3/, both sexes	4.1	11.5	35.1
Male	5.4	13.4	38.6
Female	3.1	10.2	33.6
White, both sexes	4.1	11.6	35.9
Male	5.3	13.3	38.8
Female	3.1	10.4	34.6
Black, both sexes	4.7	10.2	26.7
Male	6.7	13.5	28.7
Female	3.3	8.2	25.8

Diseases of appendix (K35-K38)

All races 3/, both sexes	0.4	1.0	1.6
Male	0.5	1.2	1.9
Female	0.3	0.8	1.5
White, both sexes	0.3	0.9	1.7
Male	0.4	1.1	1.9
Female	0.3	0.7	1.6
Black, both sexes	*	*	*
Male	*	*	*
Female	*	*	*

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

65-74 YEARS	75-84 YEARS	85 YEARS AND OVER
----------------	----------------	----------------------

Hernia (K40-K46)

All races 3/, both sexes	1.2	3.8	12.9
Male	1.1	3.9	11.9
Female	1.2	3.7	13.4
White, both sexes	1.2	3.8	13.2
Male	1.1	4.0	12.1
Female	1.2	3.6	13.6
Black, both sexes	1.2	4.2	12.5
Male	*	*	*
Female	*	4.6	12.9

Chronic liver disease and cirrhosis (K70,K73-K74)

All races 3/, both sexes	30.2	31.1	22.8
Male	40.4	37.8	32.6
Female	21.7	26.6	18.6
White, both sexes	31.0	32.3	23.7
Male	41.6	39.2	33.9
Female	22.2	27.7	19.3
Black, both sexes	25.6	18.3	13.8
Male	36.1	23.8	*
Female	17.9	15.0	12.0

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

65-74 75-84 85 YEARS
YEARS YEARS AND OVER

Alcoholic liver disease (K70)

All races 3/, both sexes	10.4	6.1	2.2
Male	17.5	11.3	5.0
Female	4.6	2.6	1.0
White, both sexes	10.6	6.1	2.3
Male	17.8	11.5	5.2
Female	4.6	2.5	1.0
Black, both sexes	10.0	5.6	*
Male	17.3	9.7	*
Female	4.7	*	*

Other chronic liver disease and cirrhosis (K73-K74)

All races 3/, both sexes	19.8	25.0	20.6
Male	23.0	26.5	27.5
Female	17.1	24.0	17.6
White, both sexes	20.4	26.2	21.4
Male	23.7	27.7	28.7
Female	17.6	25.2	18.2
Black, both sexes	15.6	12.8	12.2
Male	18.7	14.1	*
Female	13.2	12.0	11.6

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

65-74 75-84 85 YEARS
YEARS YEARS AND OVER

Cholelithiasis and other disorders of gallbladder (K80-K82)

All races 3/, both sexes	2.5	7.5	24.5
Male	2.9	9.3	26.2
Female	2.1	6.3	23.8
White, both sexes	2.3	7.4	25.3
Male	2.8	9.4	27.5
Female	1.9	6.1	24.4
Black, both sexes	3.8	8.4	17.9
Male	3.5	7.9	*
Female	4.1	8.8	20.0

Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)

All races 3/, both sexes	38.5	101.2	274.2
Male	43.4	125.2	361.2
Female	34.4	85.0	236.7
White, both sexes	32.4	94.3	265.7
Male	38.3	119.1	358.8
Female	27.5	77.4	225.9
Black, both sexes	97.4	198.8	407.4
Male	100.1	224.9	452.3
Female	95.4	182.9	388.6

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

65-74 75-84 85 YEARS
YEARS YEARS AND OVER

Acute and rapidly progressive nephritic and nephrotic syndrome (N00-N01,N04)

All races 3/, both sexes	0.2	0.4	1.1
Male	*	0.5	1.7
Female	*	0.4	0.9
White, both sexes	0.1	0.4	1.1
Male	*	0.5	*
Female	*	0.3	1.0
Black, both sexes	*	*	*
Male	*	*	*
Female	*	*	*

Chronic glomerulonephritis, nephritis and nephropathy not specified as acute or chronic, and renal sclerosis unspecified (N02-N03, N05-N07,N26)

All races 3/, both sexes	0.6	1.4	2.7
Male	0.7	1.7	4.3
Female	0.5	1.2	2.1
White, both sexes	0.5	1.4	2.4
Male	0.6	1.7	3.8
Female	0.5	1.1	1.8
Black, both sexes	*	*	*
Male	*	*	*
Female	*	*	*

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

65-74 75-84 85 YEARS
YEARS YEARS AND OVER

Renal failure (N17-N19)

All races 3/, both sexes	37.7	99.3	270.1
Male	42.6	122.9	355.0
Female	33.7	83.3	233.5
White, both sexes	31.7	92.4	262.0
Male	37.4	116.9	353.3
Female	26.8	75.8	223.0
Black, both sexes	96.1	196.0	401.4
Male	98.7	222.6	443.8
Female	94.3	179.8	383.7

Other disorders of kidney (N25,N27)

All races 3/, both sexes	*	*	*
Male	*	*	*
Female	*	*	*
White, both sexes	*	*	*
Male	*	*	*
Female	*	*	*
Black, both sexes	*	*	*
Male	*	*	*
Female	*	*	*

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

65-74 YEARS	75-84 YEARS	85 YEARS AND OVER
----------------	----------------	----------------------

Infections of kidney (N10-N12,N13.6,N15.1)

All races 3/, both sexes	0.6	2.0	7.4
Male	0.3	1.5	5.3
Female	0.8	2.4	8.4
White, both sexes	0.5	1.9	7.5
Male	*	1.4	5.6
Female	0.8	2.2	8.3
Black, both sexes	*	3.1	7.8
Male	*	*	*
Female	*	3.8	9.3

Hyperplasia of prostate (N40)

All races 3/, both sexes	0.2	1.3	5.2
Male	0.5	3.2	17.2
White, both sexes	0.2	1.3	5.2
Male	0.5	3.2	17.3
Black, both sexes	*	*	*
Male	*	*	*

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

65-74 YEARS	75-84 YEARS	85 YEARS AND OVER
----------------	----------------	----------------------

Inflammatory diseases of female pelvic organs (N70-N76)

All races 3/, both sexes	0.1	0.2	0.6
Female	0.2	0.4	0.9
White, both sexes	*	0.2	0.6
Female	*	0.4	0.9
Black, both sexes	*	*	*
Female	*	*	*

Pregnancy, childbirth and the puerperium (000-099)

All races 3/, both sexes
Female
White, both sexes
Female
Black, both sexes
Female

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

65-74 YEARS	75-84 YEARS	85 YEARS AND OVER
----------------	----------------	----------------------

Pregnancy with abortive outcome (000-007)

All races 3/, both sexes
Female
White, both sexes
Female
Black, both sexes
Female

Other complications of pregnancy, childbirth and the puerperium (010-099)

All races 3/, both sexes
Female
White, both sexes
Female
Black, both sexes
Female

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

65-74 75-84 85 YEARS
YEARS YEARS AND OVER

Certain conditions originating in the perinatal period (P00-P96)

All races 3/, both sexes	*	*	*
Male	*	*	*
Female	*	*	*
White, both sexes	*	*	*
Male	*	*	*
Female	*	*	*
Black, both sexes	*	*	*
Male	*	*	*
Female	*	*	*

Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)

All races 3/, both sexes	2.6	4.0	7.4
Male	2.7	4.0	10.1
Female	2.4	4.0	6.3
White, both sexes	2.6	4.1	7.8
Male	2.6	4.2	10.6
Female	2.6	4.1	6.5
Black, both sexes	2.5	2.8	*
Male	3.6	*	*
Female	*	*	*

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

65-74 75-84 85 YEARS
YEARS YEARS AND OVER

Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

All races 3/, both sexes	16.7	40.8	205.4
Male	20.8	45.2	173.4
Female	13.4	37.9	219.2
White, both sexes	15.9	40.3	209.9
Male	19.5	44.7	178.2
Female	12.8	37.3	223.4
Black, both sexes	28.2	56.5	186.9
Male	38.2	63.2	149.0
Female	21.0	52.5	202.8

All other diseases (Residual) (D65-E07, E15-E34, E65-F99, G04-G12, G23-G25, G31-H93, K00-K22, K29-K31, K50-K66, K71-K72, K75-K76, K83-M99, N13.0-N13.5, N13.7-N13.9, N14, N15.0, N15.8-N15.9, N20-N23, N28-N39, N41-N64, N80-N98)

All races 3/, both sexes	137.6	390.0	1,426.8
Male	150.2	384.9	1,201.4
Female	127.3	393.4	1,523.9
White, both sexes	134.6	387.9	1,448.3
Male	145.6	380.7	1,215.9
Female	125.3	392.8	1,547.7
Black, both sexes	187.1	476.4	1,392.5
Male	218.3	508.4	1,222.8
Female	164.4	457.0	1,463.4

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

65-74 YEARS	75-84 YEARS	85 YEARS AND OVER
----------------	----------------	----------------------

Accidents (unintentional injuries) (V01-X59,Y85-Y86)

All races 3/, both sexes	42.4	95.5	269.9
Male	56.5	122.2	341.2
Female	30.7	77.5	239.2
White, both sexes	41.7	96.7	279.6
Male	55.0	122.2	352.7
Female	30.6	79.4	248.4
Black, both sexes	51.1	86.4	181.6
Male	75.9	128.8	254.3
Female	33.1	60.7	151.2

Transport accidents (V01-V99,Y85)

All races 3/, both sexes	18.2	27.1	31.1
Male	24.6	37.4	56.8
Female	13.0	20.2	20.0
White, both sexes	18.1	27.6	31.9
Male	24.1	37.5	58.8
Female	13.1	20.8	20.3
Black, both sexes	19.1	20.3	21.6
Male	30.8	35.3	41.5
Female	10.7	11.3	13.3

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

65-74 75-84 85 YEARS
YEARS YEARS AND OVER

Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)

All races 3/, both sexes	16.7	25.8	30.0
Male	22.0	35.0	54.6
Female	12.4	19.5	19.4
White, both sexes	16.6	26.2	30.7
Male	21.5	35.0	56.4
Female	12.5	20.1	19.8
Black, both sexes	17.7	19.2	20.4
Male	28.1	33.5	40.4
Female	10.2	10.5	12.0

Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9,V10-V11,V15-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.6-V80.9,V81.2-V81.9,V82.2-V82.9,V87.9,V88.9,V89.1,V89.3,V89.9)

All races 3/, both sexes	0.6	0.7	0.8
Male	0.9	1.1	*
Female	0.3	0.5	*
White, both sexes	0.5	0.7	0.8
Male	0.9	1.2	*
Female	0.3	0.4	*
Black, both sexes	*	*	*
Male	*	*	*
Female	*	*	*

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

65-74 75-84 85 YEARS
YEARS YEARS AND OVER

Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)

All races 3/, both sexes	1.0	0.6	*
Male	1.8	1.2	*
Female	0.3	*	*
White, both sexes	1.0	0.7	*
Male	1.8	1.3	*
Female	0.3	*	*
Black, both sexes	*	*	*
Male	*	*	*
Female	*	*	*

Nontransport accidents (W00-X59,Y86)

All races 3/, both sexes	24.1	68.3	238.8
Male	31.9	84.7	284.5
Female	17.7	57.3	219.2
White, both sexes	23.6	69.2	247.8
Male	30.9	84.7	294.0
Female	17.4	58.6	228.0
Black, both sexes	32.0	66.1	159.9
Male	45.2	93.5	212.8
Female	22.4	49.5	137.8

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

65-74 75-84 85 YEARS
YEARS YEARS AND OVER

Falls (W00-W19)

All races 3/, both sexes	9.1	31.2	111.1
Male	12.1	39.2	138.0
Female	6.7	25.8	99.5
White, both sexes	9.3	32.7	117.8
Male	12.3	40.7	145.7
Female	6.8	27.2	105.8
Black, both sexes	8.1	15.2	38.9
Male	10.9	20.6	56.4
Female	6.0	12.0	31.6

Accidental discharge of firearms (W32-W34)

All races 3/, both sexes	0.2	0.2	*
Male	0.4	0.5	*
Female	*	*	*
White, both sexes	0.2	0.2	*
Male	0.4	0.5	*
Female	*	*	*
Black, both sexes	*	*	*
Male	*	*	*
Female	*	*	*

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

65-74 YEARS	75-84 YEARS	85 YEARS AND OVER
----------------	----------------	----------------------

Accidental drowning and submersion (W65-W74)

All races 3/, both sexes	1.0	1.3	1.5
Male	1.6	1.8	2.6
Female	0.5	0.9	1.0
White, both sexes	1.0	1.2	1.4
Male	1.6	1.8	2.6
Female	0.5	0.8	1.0
Black, both sexes	*	*	*
Male	*	*	*
Female	*	*	*

Accidental exposure to smoke, fire and flames (X00-X09)

All races 3/, both sexes	2.2	3.8	6.1
Male	2.8	4.9	8.9
Female	1.7	3.1	4.9
White, both sexes	1.9	3.1	5.1
Male	2.3	4.0	7.8
Female	1.5	2.5	3.9
Black, both sexes	5.8	13.4	20.4
Male	8.4	17.9	25.5
Female	3.9	10.7	18.2

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

65-74 75-84 85 YEARS
YEARS YEARS AND OVER

Accidental poisoning and exposure to noxious substances (X40-X49)

All races 3/, both sexes	1.5	2.0	4.3
Male	1.8	2.1	5.9
Female	1.3	1.9	3.6
White, both sexes	1.5	2.1	4.0
Male	1.6	2.1	5.6
Female	1.4	2.0	3.4
Black, both sexes	2.5	*	8.2
Male	4.3	*	*
Female	*	*	*

Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)

All races 3/, both sexes	10.0	29.9	115.6
Male	13.2	36.3	128.3
Female	7.5	25.5	110.1
White, both sexes	9.7	29.9	119.2
Male	12.6	35.6	131.3
Female	7.2	26.0	114.0
Black, both sexes	14.6	34.3	90.3
Male	19.4	51.5	114.9
Female	11.0	23.9	80.0

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

65-74 YEARS	75-84 YEARS	85 YEARS AND OVER
----------------	----------------	----------------------

Intentional self-harm (suicide) (X60-X84,Y87.0)

All races 3/, both sexes	12.6	17.7	19.4
Male	23.0	38.0	54.5
Female	4.1	4.0	4.3
White, both sexes	13.7	18.8	20.8
Male	24.7	40.4	58.8
Female	4.4	4.1	4.5
Black, both sexes	5.0	5.9	*
Male	10.6	11.8	*
Female	*	*	*

Intentional self-harm (suicide) by discharge of firearms (X72-X74)

All races 3/, both sexes	9.4	13.1	13.0
Male	18.6	30.3	40.7
Female	1.7	1.4	1.1
White, both sexes	10.2	14.2	14.1
Male	20.1	32.8	44.3
Female	2.0	1.5	1.2
Black, both sexes	3.7	3.6	*
Male	8.4	7.9	*
Female	*	*	*

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

65-74 75-84 85 YEARS
YEARS YEARS AND OVER

Intentional self-harm (suicide) by other and unspecified means and their sequelae (X60-X71,X75-X84,Y87.0)

All races 3/, both sexes	3.3	4.6	6.4
Male	4.4	7.6	13.8
Female	2.3	2.6	3.2
White, both sexes	3.4	4.7	6.7
Male	4.5	7.7	14.5
Female	2.5	2.6	3.3
Black, both sexes	1.4	2.3	*
Male	*	*	*
Female	*	*	*

Assault (homicide) (X85-Y09,Y87.1)

All races 3/, both sexes	2.4	2.4	2.3
Male	3.4	3.1	3.1
Female	1.6	2.0	2.0
White, both sexes	1.9	2.0	2.1
Male	2.4	2.4	2.7
Female	1.5	1.8	1.9
Black, both sexes	7.4	7.1	*
Male	13.3	11.5	*
Female	3.1	4.5	*

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

65-74 75-84 85 YEARS
YEARS YEARS AND OVER

Assault (homicide) by discharge of firearms (X93-X95)

All races 3/, both sexes	1.0	0.7	0.7
Male	1.3	0.7	*
Female	0.8	0.6	*
White, both sexes	0.8	0.6	0.6
Male	1.0	0.6	*
Female	0.7	0.6	*
Black, both sexes	2.7	*	*
Male	5.2	*	*
Female	*	*	*

Assault (homicide) by other and unspecified means and their sequelae (X85-X92,X96-Y09,Y87.1)

All races 3/, both sexes	1.4	1.8	1.7
Male	2.1	2.4	2.4
Female	0.9	1.4	1.4
White, both sexes	1.1	1.5	1.5
Male	1.5	1.9	2.0
Female	0.7	1.2	1.3
Black, both sexes	4.7	5.2	*
Male	8.1	7.9	*
Female	2.2	3.6	*

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

65-74 75-84 85 YEARS
YEARS YEARS AND OVER

Legal intervention (Y35,Y89.0)

All races 3/, both sexes	*	*	*
Male	*	*	*
Female	*	*	*
White, both sexes	*	*	*
Male	*	*	*
Female	*	*	*
Black, both sexes	*	*	*
Male	*	*	*
Female	*	*	*

Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)

All races 3/, both sexes	0.6	0.6	1.3
Male	0.9	1.0	1.6
Female	0.3	0.4	1.2
White, both sexes	0.5	0.6	1.2
Male	0.7	1.0	*
Female	0.4	0.4	1.1
Black, both sexes	*	*	*
Male	*	*	*
Female	*	*	*

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

65-74 75-84 85 YEARS
YEARS YEARS AND OVER

Discharge of firearms, undetermined intent (Y22-Y24)

All races 3/, both sexes	*	*	*
Male	*	*	*
Female	*	*	*
White, both sexes	*	*	*
Male	*	*	*
Female	*	*	*
Black, both sexes	*	*	*
Male	*	*	*
Female	*	*	*

Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)

All races 3/, both sexes	0.5	0.6	1.3
Male	0.7	0.9	*
Female	0.3	0.4	1.2
White, both sexes	0.5	0.6	1.2
Male	0.6	0.9	*
Female	0.3	0.4	1.1
Black, both sexes	*	*	*
Male	*	*	*
Female	*	*	*

TABLE 250. DEATH RATES FOR 113 SELECTED CAUSES, BY 10-YEAR AGE GROUPS, RACE, AND SEX: UNITED STATES, 2000

(RATES PER 100,000 POPULATION IN SPECIFIED GROUP. POPULATIONS USED FOR COMPUTING DEATH RATES ARE POSTCENSAL ESTIMATES BASED ON THE 1990 CENSUS, ESTIMATED AS OF JULY 1, 2000)

CAUSE OF DEATH (BASED ON
THE TENTH REVISION, INTER-
NATIONAL CLASSIFICATION
OF DISEASES, 1992),
RACE, AND SEX

65-74 75-84 85 YEARS
YEARS YEARS AND OVER

Operations of war and their sequelae (Y36,Y89.1)

All races 3/, both sexes	*	*	*
Male	*	*	*
Female	*	*	*
White, both sexes	*	*	*
Male	*	*	*
Female	*	*	*
Black, both sexes	*	*	*
Male	*	*	*
Female	*	*	*

Complications of medical and surgical care (Y40-Y84,Y88)

All races 3/, both sexes	3.6	6.7	13.5
Male	4.0	8.1	15.2
Female	3.2	5.8	12.7
White, both sexes	3.4	6.5	13.0
Male	3.8	8.1	15.1
Female	3.0	5.4	12.1
Black, both sexes	5.9	10.7	21.0
Male	7.3	10.0	*
Female	5.0	11.1	22.7

* FIGURE DOES NOT MEET STANDARDS OF RELIABILITY OR PRECISION.

0.0 QUANTITY MORE THAN ZERO BUT LESS THAN 0.05.

... CATEGORY NOT APPLICABLE.

1/ FIGURES FOR AGE NOT STATED INCLUDED IN "ALL AGES" BUT NOT DISTRIBUTED AMONG AGE GROUPS.

2/ DEATH RATES FOR "UNDER 1 YEAR" (BASED ON POPULATION ESTIMATES) DIFFER FROM INFANT MORTALITY RATES (BASED ON LIVE BIRTHS).

3/ INCLUDES RACES OTHER THAN WHITE AND BLACK.

SOURCE: CDC/NCHS, NATIONAL VITAL STATISTICS SYSTEM, MORTALITY.